

# Vertex-Transitive Cubic Graphs of Square-Free Order\*

---

Cai Heng Li,<sup>1,2</sup> Zai Ping Lu,<sup>3</sup> and Gai Xia Wang<sup>4</sup>

<sup>1</sup>SCHOOL OF MATHEMATICS AND STATISTICS  
YUNNAN UNIVERSITY, KUNMING, YUNNAN 650091, P. R. CHINA

<sup>2</sup>SCHOOL OF MATHEMATICS AND STATISTICS  
THE UNIVERSITY OF WESTERN AUSTRALIA, CRAWLEY, WA 6009, AUSTRALIA  
E-mail: cai.heng.li@uwa.edu.au

<sup>3</sup>CENTER FOR COMBINATORICS  
LPMC-TJKLC, NANKAI UNIVERSITY, TIANJIN 300071, P. R. CHINA  
E-mail: lu@nankai.edu.cn

<sup>4</sup>DEPARTMENT OF APPLIED MATHEMATICS  
ANHUI UNIVERSITY OF TECHNOLOGY, MAANSHAN 243002, P. R. CHINA  
E-mail: gxwang@ahut.edu.cn

Received November 5, 2011; Revised September 16, 2012

Published online 2 January 2013 in Wiley Online Library (wileyonlinelibrary.com).  
DOI 10.1002/jgt.21715

**Abstract:** A classification of connected vertex-transitive cubic graphs of square-free order is provided. It is shown that such graphs are well-characterized metacirculants (including dihedrants, generalized Petersen graphs, Möbius bands), or Tutte's 8-cage, or graphs arisen from simple groups  $\text{PSL}(2, p)$ . © 2012 Wiley Periodicals, Inc. *J. Graph Theory* 75: 1–19, 2014

**Keywords:** vertex-transitive graph; arc-transitive graph; metacirculant; Tutte's 8-cage; coset graph

---

\*This work was partially supported by two Natural Science Funds of China, a Tianyuan Youth Fund of China, a Yunnan High-End-Talent-Plan fund, and an ARC Discovery Project grant of Australia.

*Journal of Graph Theory*  
© 2012 Wiley Periodicals, Inc.

## 1. INTRODUCTION

For a graph  $\Gamma = (V, E)$ , the number of vertices  $|V|$  is called the *order* of  $\Gamma$ . A graph  $\Gamma$  is called *vertex-transitive* if its automorphism group  $\text{Aut}\Gamma$  is transitive on  $V$ .

In 1967, Turner [22] investigated vertex-transitive graphs of prime order, and enumerated the isomorphism classes of such graphs by using Pólya enumeration theorem. Since then, the class of vertex-transitive graphs of square-free order has been studied extensively and numerous interesting results have appeared on classification, isomorphism problem, non-Cayley numbers, etc. Classification results about vertex-transitive graphs of square-free order usually focus on specific subclasses regarding their symmetry properties, orders, valencies, etc. For instance, see [18, 20] for those graphs of order being a product of two prime, see [1, 4, 5, 9, 10, 15, 17, 19, 24] for those graphs having certain symmetry properties. In a recent paper [23], a classification was given of vertex-transitive cubic graphs of order  $2pq$ , where  $p$  and  $q$  are primes.

In this article, we classify vertex-transitive cubic graphs of square-free order.

A graph is called a *metacirculant* if it has a vertex-transitive metacyclic group of automorphisms. Examples of vertex-transitive cubic graphs of square-free order include a lot of interesting graphs:  $K_{3,3}$ , Petersen graph, Tutte's 8-cage (30 vertices), generalized Petersen graphs, Möbius bands, some well-characterized metacirculants, and some graphs arisen from simple groups  $\text{PSL}(2, p)$ . See Section 2 for definitions and constructions. Among these graphs, some are Cayley graphs. For a group  $G$  and a subset  $S \subset G$  with  $1 \notin S = S^{-1} := \{g^{-1} \mid g \in S\}$ , the *Cayley graph*  $\text{Cay}(G, S)$  is defined on  $G$  such that  $\{g, h\}$  is an edge if and only if  $gh^{-1} \in S$ .

Throughout this article, for two groups  $A$  and  $B$ , denote by  $A \times B$ ,  $A.B$  and  $A:B$  the direct product, an extension and a semidirect product of  $A$  by  $B$ , respectively; denote, respectively, by  $A'$  and  $\mathbf{Z}(A)$  the commutator subgroup and the center of  $A$ ; for  $a \in A$ , denote by  $o(a)$  the order of  $a$  in  $A$ ; for a positive integer  $n$ , denote by  $\mathbb{Z}_n$  and  $D_{2n}$  the cyclic group of order  $n$  and the dihedral group of order  $2n$ , respectively.

Our classification is stated in the following theorem.

**Theorem 1.1.** *Let  $\Gamma$  be a connected vertex-transitive cubic graph of square-free order  $2n$ . Then one of the following statements holds.*

- (1)  $\Gamma$  is a metacirculant, and one of the following is true:
  - (i)  $\Gamma$  is isomorphic to a generalized Petersen graph  $\mathbf{P}(n, r)$  for  $1 \leq r < \frac{n}{2}$  with  $r^2 \equiv 1 \pmod{n}$ ;  $\text{Aut}\Gamma \cong \mathbb{Z}_n : \mathbb{Z}_2^2$  has a regular subgroup  $\langle a, b \mid a^n = b^2 = 1, bab = a^r \rangle$ , and has no regular subgroups isomorphic to  $\mathbb{Z}_{2n}$  or  $D_{2n}$  unless  $r = 1$ ;
  - (ii)  $\Gamma$  is the Möbius band  $\mathbf{M}_n$  of order  $2n$ ; either  $\text{Aut}\Gamma \cong \mathbb{Z}_{2n} : \mathbb{Z}_2 \cong D_{4n}$  or  $\Gamma \cong K_{3,3}$ ;
  - (iii)  $\Gamma \cong \text{Cay}(\langle a, b \rangle, S)$  for  $S = \{ab, a^k b, b\}$  or  $\{ab, a^{1-k} b, b\}$ ,  $\langle a, b \rangle \cong D_{2n}$ ,  $o(a) = n > 3$  and  $o(b) = 2$ , where  $k \not\equiv -1 \pmod{n}$  and  $k^2 \equiv 1 \pmod{n}$ ; in this case,  $\text{Aut}\Gamma \cong D_{2n} : \mathbb{Z}_2$  contains no cyclic regular subgroups;
  - (iv)  $\Gamma \cong \text{Cay}(\langle a, b \rangle, \{ab, a^k b, b\})$  for  $\langle a, b \rangle \cong D_{2n}$ ,  $o(a) = n > 3$  and  $o(b) = 2$ , where  $k^2 - k + 1 \equiv 0 \pmod{n}$ ; in this case,  $\text{Aut}\Gamma \cong D_{2n} : \mathbb{Z}_3$  except for Line 1 of Table I;
  - (v)  $\Gamma \cong \text{Cay}(\langle a, b \rangle, \{a^{k'} b, a^k b, b\})$  for  $\langle a, b \rangle \cong D_{2n}$ ,  $o(a) = n > 3$  and  $o(b) = 2$ , where  $(k, k') = 1$ , either  $(k, n) \neq 1$  and  $(k', n) \neq 1$ , or  $k' \equiv 1 \pmod{n}$ ,

TABLE I. Some exceptions.

Line	Regular subgroup	$k$	$\text{Aut}\Gamma$	$\Gamma (\cong)$
1	$\langle a, b \rangle \cong D_{14}$	3 or 5	$\text{PGL}(2, 7)$	Example 3.6 (2)
2	$\langle a, b \rangle \cong \mathbb{Z}_7:\mathbb{Z}_6$	2	$\text{PGL}(2, 7)$	Example 3.8 (2)
3	$\langle a, b \rangle \cong \mathbb{Z}_{\frac{n}{3}}:\mathbb{Z}_6, a^b = a^t$ $t^2 - t + 1 \equiv 0 \pmod{n}$	1	$D_{2n}:\mathbb{Z}_3$	Lemma 2.3 (3)
4	$\langle a, b \rangle \cong \mathbb{Z}_{11}:\mathbb{Z}_{10}$	$a^{b^k} = a^7 \text{ or } a^8$	$\text{PGL}(2, 11)$	Example 3.6 (1)
5	$\langle a, b \rangle \cong \mathbb{Z}_{23}:\mathbb{Z}_{22}$	$a^{b^k} = a^{17} \text{ or } a^{19}$	$\text{PGL}(2, 23)$	Example 3.6 (2)

- $k^2 \not\equiv 1 \pmod{n}$ ,  $(k-1)^2 \not\equiv 1 \pmod{n}$ ,  $2k \not\equiv 1 \pmod{n}$ , and  $k^2 - k + 1 \not\equiv 0 \pmod{n}$ ; in this case,  $\text{Aut}\Gamma \cong \langle a, b \rangle$ ;  
 (vi)  $\Gamma \cong \text{Cay}(\langle a, b, c \rangle, \{cab^k, (cab^k)^{-1}, b^l\})$ ,  $\mathbf{Z}(\langle a, b, c \rangle) = \langle c \rangle$ ,  $(\langle a, b, c \rangle)' = \langle a \rangle$ ,  $2 < o(a) < n$ ,  $2 < o(b) = 2l$  and  $a^{b^l} = a^{-1}$ , where  $0 < k < l$  and  $(k, l) = 1$ ; in this case,  $\text{Aut}\Gamma \cong \langle a, b, c \rangle$  except for Lines 2–5 of Table I;  
 (vii)  $\Gamma \cong \mathbf{P}(n, r)$  with  $1 < r < \frac{n}{2}$  and  $r^2 \equiv -1 \pmod{n}$ ; either  $\text{Aut}\Gamma \cong \mathbb{Z}_n:\mathbb{Z}_4$ , or  $\text{Aut}\Gamma = S_5$  and  $\Gamma$  is isomorphic to the Petersen graph;

- (2)  $\Gamma$  is isomorphic to Tutte's 8-cage,  $n = 15$  and  $\text{Aut}\Gamma = \text{P}\Gamma\text{L}(2, 9)$ ;  
 (3)  $\text{Aut}\Gamma = \text{PSL}(2, p)$  or  $\text{PGL}(2, p)$  for a prime  $p \geq 5$ , and  $\Gamma$  is isomorphic to one of the graphs constructed in Examples 3.5–3.8;  
 (4)  $\text{Aut}\Gamma = \text{PSL}(2, p):D_{2m}$  for a prime  $p \geq 5$  and  $1 < m = \frac{8n}{p(p^2-1)}$ , and  $\Gamma$  is isomorphic to one of the graphs constructed in Construction 4.2.

We remark that a characterization of general cubic metacirculants was given in [16], in which two families of such graphs are proved to be covers of some special graphs but the covers are not yet determined. Part (1) of Theorem 1.1 gives an explicit classification of cubic metacirculants of square-free order.

## 2. CUBIC METACIRCULANTS

Let  $n \geq 3$  and  $1 \leq r < \frac{n}{2}$  be two integers. The *generalized Petersen graph*  $\mathbf{P}(n, r)$  is the graph with vertex set and edge set as follows

$$\begin{aligned}
 &\{\alpha_0, \alpha_1, \dots, \alpha_{n-1}\} \cup \{\beta_0, \beta_1, \dots, \beta_{n-1}\}, \\
 &\{\{\alpha_i, \alpha_{i+1}\}, \{\alpha_i, \beta_i\}, \{\beta_i, \beta_{i+r}\} \mid 0 \leq i \leq n-1\},
 \end{aligned}$$

reading  $i+1$  and  $i+r$  modulo  $n$ . It was shown in [11] that  $\mathbf{P}(n, r)$  is vertex-transitive if and only if either  $(n, r) = (10, 2)$  or  $r^2 \equiv \pm 1 \pmod{n}$ . Further,  $\text{Aut}\mathbf{P}(n, r)$  has a transitive subgroup isomorphic to  $\mathbb{Z}_n:\mathbb{Z}_4$  if  $r^2 \equiv -1 \pmod{n}$ , and has a regular subgroup isomorphic to  $\mathbb{Z}_n:\mathbb{Z}_2$  if  $r^2 \equiv 1 \pmod{n}$ . In particular,  $\text{Aut}\mathbf{P}(n, 1)$  contains two regular subgroups isomorphic to  $\mathbb{Z}_{2n}$  and  $D_{2n}$ , respectively.

The *Möbius band*  $\mathbf{M}_n$  of order  $2n$  is the graph with vertex set  $\{\alpha_0, \alpha_1, \dots, \alpha_{2n-1}\}$ , and edge set  $\{\{\alpha_i, \alpha_{i+1}\}, \{\alpha_i, \alpha_{i+n}\} \mid 0 \leq i \leq 2n-1\}$ , reading the subscripts modulo  $2n$ . For the graph  $\mathbf{M}_n$ , its automorphism group contains two regular subgroups isomorphic to  $\mathbb{Z}_{2n}$  and  $D_{2n}$ , respectively.

A graph  $\Gamma = (V, E)$  is called a *circulant* or *dihedrant* if  $\text{Aut}\Gamma$  contains, respectively, a cyclic or dihedral subgroup which is regular on the vertex set  $V$ .

Let  $\Gamma = (V, E)$  be a graph such that  $\text{Aut}\Gamma$  has a regular subgroup  $G$ . Take  $\alpha \in V$ . Then each vertex of  $\Gamma$  is uniquely written as  $\alpha^g$  for some  $g \in G$ . Let  $\Gamma(\alpha)$  be the set of neighbors of  $\alpha$  in  $\Gamma$ . Set  $S = \{s \in G \mid \alpha^s \in \Gamma(\alpha)\}$ . Then  $1 \notin S = S^{-1} := \{s^{-1} \mid s \in S\}$  and  $\Gamma \cong \text{Cay}(G, S)$ . It is well known that a Cayley graph  $\text{Cay}(G, S)$  is connected whenever  $S$  generates the underlying group  $G$ , that is,  $\langle S \rangle = G$ . Moreover, each automorphism  $\sigma \in \text{Aut}(G)$  of the group  $G$  induces naturally an isomorphism from  $\text{Cay}(G, S)$  to  $\text{Cay}(G, S^\sigma)$ . Set

$$\text{Aut}(G, S) = \{\sigma \in \text{Aut}(G) \mid S^\sigma = S\}.$$

For  $g \in G$ , by  $\bar{g}$  we denote the permutation induced by  $g$  on  $G$  by right multiplication. Set  $\bar{G} = \{\bar{g} \mid g \in G\}$ . Then  $G \rightarrow \bar{G}, g \mapsto \bar{g}$  is an isomorphism of groups. By [12, Lemma 2.1], the normalizer  $\mathbf{N}_{\text{AutCay}(G, S)}(\bar{G}) = \bar{G} : \text{Aut}(G, S)$ .

To end this section, let  $G$  be a group of square-free order  $2n$ . Then  $n$  is odd.

**Lemma 2.1.** *For a group  $G$  of square-free order  $2n$ , one of the following holds.*

- (1)  $G \cong \mathbb{Z}_{2n}$  or  $\text{D}_{2n}$ ;
- (2)  $G' \cong \mathbb{Z}_m$  and  $G \cong \mathbb{Z}_m : \mathbb{Z}_{\frac{2n}{m}}$  for odd  $m$  with  $n > m > 2$ .

**Proof.** Since  $G$  has square-free order,  $G'$  is cyclic and  $G = G' : H$ , where  $H$  is a cyclic Hall subgroup of  $G$ . Set  $G' = \langle a \rangle$  and  $H = \langle b \rangle$ . If  $G' = 1$ , then  $G = H \cong \mathbb{Z}_{2n}$ .

Let  $G' = \langle a \rangle \cong \mathbb{Z}_m$  for  $m > 1$ . If  $m$  is even, then  $a^{\frac{m}{2}}$  lies in the center of  $G$ , so  $G/\langle a^2 \rangle \cong \langle a^{\frac{m}{2}}, b \rangle$  is abelian, hence  $G' = \langle a \rangle \leq \langle a^2 \rangle$ , which is impossible. Thus  $m$  is odd, and so  $H$  is of even order  $\frac{2n}{m}$ . If  $n > m$ , then part (2) occurs. Assume that  $m = n$ . Let  $C = \mathbf{C}_{\langle a \rangle}(b)$ . Then there is a subgroup  $D$  of  $\langle a \rangle$  with  $\langle a \rangle = C \times D$ . It is easily shown that  $D$  is normal in  $G$ . Then  $G/D \cong C \times \langle b \rangle$  is abelian, so  $G' \leq D$ , hence  $D = \langle a \rangle$  and  $C = 1$ . It follows that  $a^b = a^{-1}$ , hence  $G \cong \text{D}_{2n}$ . ■

Let  $\Gamma \cong \text{Cay}(G, S)$ , where  $S$  be a generating set of  $G$  with  $|S| = 3$  and  $1 \notin S = S^{-1}$ . Then  $S$  either contains only one involution, or consists of involutions. Since  $\Gamma$  is connected,  $\langle S \rangle = G$ , we know that  $\text{Aut}(G, S)$  is faithful on  $S$ . It follows that  $\text{Aut}(G, S)$  is isomorphic to a subgroup of the symmetric group  $\text{S}_3$  of degree 3.

Let  $G$  be abelian. Then  $G$  is cyclic,  $S = \{x, x^{-1}, z\}$  and  $\text{Aut}(G, S) \cong \mathbb{Z}_2$ , where  $z$  is the unique involution in  $G$ . Since  $\langle S \rangle = G$ , either  $G = \langle x \rangle$  or  $G = \langle x \rangle \times \langle z \rangle$ . If  $G = \langle x \rangle \times \langle z \rangle$ , then  $\Gamma \cong \mathbf{P}(n, 1)$ . Let  $G = \langle x \rangle$ . Then  $z = x^n$ . Set  $\alpha_i = x^i$ . Then  $\alpha_i$  and  $\alpha_j$  are adjacent whenever  $j - i \equiv \pm 1 \pmod{2n}$  or  $j - i \equiv n \pmod{2n}$ . Thus  $\Gamma \cong \mathbf{M}_n$ , and the next result follows.

**Lemma 2.2.** *A connected cubic circulant of order  $2n$  is either the ladder graph  $\mathbf{P}(n, 1)$  or the Möbius band  $\mathbf{M}_n$ .*

Thus we assume next that  $G$  is not abelian. Since  $G$  has square-free order, a Sylow 2-subgroup of  $G$  has order 2, it follows that all involutions in  $G$  are conjugate. The next lemma give a characterization of connected cubic dihedrants.

**Lemma 2.3.** *Let  $G$  the dihedral group of order  $2n$ , and let  $\Gamma$  be a connected cubic Cayley graph of  $G$ . Set  $G = \langle a, b \rangle$  with  $o(a) = n$ ,  $o(b) = 2$ , and  $a^b = a^{-1}$ . Then  $\Gamma \cong \text{Cay}(G, S)$  for one of the following subset  $S$  of  $G$ .*

- (1)  $S = \{a, a^{-1}, b\}$ ; in this case,  $\text{Aut}(G, S) \cong \mathbb{Z}_2$  and  $\Gamma \cong \mathbf{P}(n, 1)$ ;
- (2)  $n = 3$  and  $S = \{ab, a^2b, b\}$ ; in this case,  $\Gamma \cong \mathbf{K}_{3,3}$ ;
- (3)  $S = \{ab, a^kb, b\}$ ,  $k^2 - k + 1 \equiv 0 \pmod{n}$ ,  $n > 3$ ; in this case,  $\text{Aut}(G, S) \cong \mathbb{Z}_3$ ;
- (4)  $S = \{ab, a^eb, b\}$  or  $\{ab, a^{1-e}b, b\}$  for  $n > 3$  and  $e^2 \equiv 1 \pmod{n}$ ; in this case,  $\text{Aut}(G, S) \cong \mathbb{Z}_2$ ;
- (5)  $S = \{ab, a^kb, b\}$ ,  $n > 3$ ,  $k^2 \not\equiv 1 \pmod{n}$ ,  $(k-1)^2 \not\equiv 1 \pmod{n}$ ,  $2k \not\equiv 1 \pmod{n}$  and  $k^2 - k + 1 \not\equiv 0 \pmod{n}$ ; in this case,  $\text{Aut}(G, S) = 1$ ;
- (6)  $S = \{a^kb, a^kb, b\}$ ,  $n > 3$ ,  $(k, k') = 1$ ,  $(k, n) \neq 1$  and  $(k', n) \neq 1$ ; in this case,  $\text{Aut}(G, S) = 1$ .

**Proof.** Let  $\Gamma = \text{Cay}(G, S)$ . Recall that all involutions in  $G$  are conjugate. Up to isomorphism of graphs we may choose  $b \in S$ . If  $S$  has only one involution, then  $S = \{a^s, a^{-s}, b\}$ , where  $(s, n) = 1$ . It is easily shown that  $\text{Aut}(G, S) \cong \mathbb{Z}_2$ . Take  $\sigma \in \text{Aut}(G)$  with  $(a^s)^\sigma = a$  and  $b^\sigma = b$ , refer to [14]. Then  $\Gamma \cong \text{Cay}(G, S^\sigma)$  and  $S^\sigma = \{a, a^{-1}, b\}$ . Set  $\alpha_i = a^i$  and  $\beta_i = ba^i$  for  $0 \leq i \leq n-1$ . Then  $\text{Cay}(G, S^\sigma)$  has edges  $\{\alpha_i, \alpha_{i+1}\}$ ,  $\{\beta_i, \beta_{i+1}\}$ , and  $\{\alpha_i, \beta_i\}$ . Thus  $\Gamma \cong \mathbf{P}(n, 1)$ .

Assume that  $S = \{x, y, b\}$  consists of three involutions. Then  $S = \{a^ib, a^jb, b\}$  for some positive integers  $i$  and  $j$ . Let  $d = (i, j)$ ,  $i = kd$ , and  $j = k'd$ . Then  $G = \langle S \rangle = \langle a^i, a^j, b \rangle = \langle a^i, a^j \rangle : \langle b \rangle = \langle a^d \rangle : \langle b \rangle$ , so  $\langle a^d \rangle = \langle a \rangle$ , hence  $(d, n) = 1$ . Thus  $sd \equiv 1 \pmod{n}$  for some  $s$  coprime to  $n$ . Take an automorphism  $\sigma \in \text{Aut}(G)$  with  $a^\sigma = a^s$  and  $b^\sigma = b$ , refer to [14]. Then  $S^\sigma = \{a^kb, a^{k'}b, b\}$  and  $\Gamma \cong \text{Cay}(G, S^\sigma)$ .

Suppose that  $\text{Aut}(G, S^\sigma)$  has an element  $\tau$  of order 3. Let  $a^\tau = a^t$  for some  $t$  coprime to  $n$ . Then  $t^3 \equiv 1 \pmod{n}$ . Noting that  $\tau^{-1} \in \text{Aut}(G, S^\sigma)$ , without loss of generality, we may set  $b^\tau = a^{k'}b$ . Since  $S^{\sigma\tau} = S^\sigma$ , computation shows that  $S^\sigma = \{b, a^{k'}b, a^{k'(t+1)}b\}$ ,  $k'(t+1) \equiv k \pmod{n}$ ,  $k'(t^2 + t + 1) \equiv 0 \pmod{n}$ . By the argument in above paragraph, we know that  $(k', n) = 1$ . Thus we have

- (i)  $S^\sigma = \{b, a^{k'}b, a^{k'(t+1)}b\}$ ,  $(k', n) = 1$ ,  $(k, n) = 1$ ,  $k'(t+1) \equiv k \pmod{n}$ ,  $(t^2 + t + 1) \equiv 0 \pmod{n}$ .

Suppose that  $\text{Aut}(G, S^\sigma)$  has an involution  $\varepsilon$ . Let  $a^\varepsilon = a^e$  for some  $e$  coprime to  $n$ . Then  $e^2 \equiv 1 \pmod{n}$ . Note that  $\varepsilon$  fixes one involution in  $S^\sigma$  and interchanges the other two. Then one of the following occurs:

- (ii)  $S^\sigma = \{a^{k'}b, a^{k'e}b, b\}$ ,  $(k', n) = 1$ ,  $(k, n) = 1$  and  $k \equiv k'e \pmod{n}$ ;
- (iii)  $S^\sigma = \{a^{k'}b, a^{k'(1-e)}b, b\}$ ,  $(k', n) = 1$ ,  $k' - k'e \equiv k \pmod{n}$ ;
- (iii)'  $S^\sigma = \{a^{(1-e)k}b, a^{k'}b, b\}$ ,  $(k, n) = 1$ ,  $k \equiv k' + ke \pmod{n}$ .

Conversely, it is easily shown that  $\text{Aut}(G, S^\sigma) \neq 1$  if  $S^\sigma$  is described as in one of the above items (i)–(iii)'. It is easily shown that  $\text{Aut}(S^\sigma) \cong S_3$  if and only if  $n = 3$ .

By the above argument,  $\text{Aut}(G, S^\sigma) = 1$  if neither  $(k, n) = 1$  nor  $(k', n) \neq 1$ , and then part (6) follows. Thus, without loss of generality, we assume next that  $(k', n) = 1$ . Then, by [14], there is  $\delta \in \text{Aut}(G)$  with  $(a^{k'})^\delta = a$  and  $b^\delta = b$ . Since  $\text{Cay}(G, S^\sigma) \cong \text{Cay}(G, S^{\sigma\delta})$ , replacing  $S^\sigma$  by  $S^{\sigma\delta}$ , we may assume that  $S^\sigma = \{ab, a^kb, b\}$ , that is, take  $k' = 1$ . If  $n = 3$ , then the part (2) of the lemma follows. Let  $n > 3$ . If item (i) holds, then part (3) follows. If item (ii) or (iii) holds, then part (4) follows. Assume that (iii)' holds then  $1 \equiv k' \equiv k(1-e) \pmod{n}$ , so  $(1-e, n) = 1$ . Hence,  $e \equiv -1 \pmod{n}$  as  $e^2 \equiv 1 \pmod{n}$ . Thus  $2k \equiv 1 \pmod{n}$ . Noting that  $(k, n) = 1$ , we may take an automorphism of  $G$  with  $a^k \mapsto a$  and  $b \mapsto b$ . Then  $\Gamma \cong \text{Cay}(G, \{ab, a^kb, b\}) \cong \text{Cay}(G, \{a^2b, ab, b\})$ , which is a

graph given in part (4). For  $S^\sigma = \{ab, a^k b, b\}$ , by the above argument,  $\text{Aut}(G, S^\sigma) = 1$  if and only if  $n > 3$ ,  $k^2 \not\equiv 1 \pmod{n}$ ,  $(k-1)^2 \not\equiv 1 \pmod{n}$ ,  $2k \not\equiv 1 \pmod{n}$ , and  $k^2 - k + 1 \not\equiv 0 \pmod{n}$ . Then part (5) follows. ■

**Corollary 2.4.** *Let  $n > 3$  and  $G = \langle a \rangle : \langle b \rangle \cong D_{2n}$  be of square-free order, and let  $S = \{ab, a^e b, b\}$  or  $\{ab, a^{1-e} b, b\}$  be as in Lemma 2.3 (4). Then  $\bar{G} : \text{Aut}(G, S)$  has a cyclic regular subgroup if and only if  $e \equiv -1 \pmod{n}$ .*

**Proof.** Let  $\Gamma = \text{Cay}(G, S)$ . Then  $\text{Aut}(G, S) = \langle \sigma \rangle \cong \mathbb{Z}_2$ , where  $\sigma \in \text{Aut}(G)$  with  $a^\sigma = a^e$  and either  $b^\sigma = b$  for  $S = \{ab, a^e b, b\}$  or  $b^\sigma = a^{1-e} b$  for  $S = \{ab, a^{1-e} b, b\}$ . Let  $g \in S$  with  $g^\sigma = g$ . It is easily shown that each regular subgroup of  $\bar{G} : \text{Aut}(G, S)$  can be written as  $R := \langle \bar{a}, \sigma^j \bar{g} \rangle$  for  $j = 0$  or  $1$ . Clearly,  $R$  is cyclic if and only if  $j = 1$  and  $\bar{a}^{-e} = \bar{a}^{\sigma \bar{g}} = (\sigma \bar{g})^{-1} \bar{a} \sigma \bar{g} = \bar{a}$ , that is,  $e \equiv -1 \pmod{n}$ . ■

Now assume that  $G$  satisfies Lemma 2.1 (2). Then  $G$  cannot be generated by three involutions. Thus, for a connected cubic graph  $\text{Cay}(G, S)$ , the subset  $S$  contains only one involution of  $G$ . Since  $G$  is not abelian, this involution is not contained in the center of  $G$ . Let  $H < G$  with  $G = G' : H$ , and let  $C = C_H(G')$ . Then  $C$  is the center of  $G$  and of odd order, and  $G = C \times (G' : \langle b \rangle)$  for a cyclic subgroup  $\langle b \rangle$  of  $H$  of even order. Set  $C = \langle c \rangle$  and  $G' = \langle a \rangle$ . Then  $o(c)o(b) > 2$ , and so  $2 < o(a) < n$ .

**Lemma 2.5.** *Let  $G = \langle c \rangle \times (\langle a \rangle : \langle b \rangle)$  be a group of square-free order  $2n$ , where  $\mathbf{Z}(G) = \langle c \rangle$  and  $G' = \langle a \rangle \cong \mathbb{Z}_m$  with  $2 < m < n$ . Let  $\Gamma$  be a connected cubic Cayley graph of  $G$ . Then  $o(b) = 2l$ ,  $a^{b^l} = a^{-1}$  and  $\Gamma \cong \text{Cay}(G, S_k)$  for  $S_k = \{cab^k, (cab^k)^{-1}, b^l\}$ , where  $l \geq 1$ ,  $0 \leq k \leq l$ , and  $(k, l) = 1$ . Moreover,  $\text{Aut}(G, S_k) \neq 1$  if and only if  $l = 1$ ; in this case, either  $\Gamma$  is a dihedrant, or  $\Gamma \cong \mathbf{P}(n, r)$  with  $1 < r < \frac{n}{2}$  and  $r^2 \equiv 1 \pmod{n}$ .*

**Proof.** Let  $\Gamma \cong \text{Cay}(G, S)$ . By the above argument,  $o(b)$  is even. Set  $o(b) = 2l$ . Recall that all involutions in  $G$  are conjugate. Up to isomorphism of graphs, we may choose  $b^l \in S$  and set  $S = \{xyz, (xyz)^{-1}, b^l\}$ , where  $x \in \langle c \rangle$ ,  $y \in \langle a \rangle$  and  $z \in \langle b \rangle$ . Since  $\langle S \rangle = G$ , we have  $\langle x \rangle = \langle c \rangle$ ,  $\langle y \rangle = \langle a \rangle$  and  $\langle z, b^l \rangle = \langle b \rangle$ . Take  $\sigma \in \text{Aut}(G)$  with  $x^\sigma = c$ ,  $y^\sigma = a$ , and  $b^\sigma = b$ , refer to [14]. Then  $S_k := S^\sigma = \{cab^k, (cab^k)^{-1}, b^l\}$  for some  $0 \leq k < 2l$  coprime to  $l$ , and so  $\Gamma \cong \text{Cay}(G, S_k)$ . Setting  $a^b = a^r$ , by [14], we may take  $\rho \in \text{Aut}(G)$  with  $c^\rho = c^{-1}$ ,  $a^\rho = a^{-r^{2l-k}}$ , and  $b^\rho = b$ . Then  $S_k^\rho = \{cab^{2l-k}, (cab^{2l-k})^{-1}, b^l\} = S_{2l-k}$ , so  $\text{Cay}(G, S_k) \cong \text{Cay}(G, S_{2l-k})$ . Thus, up to isomorphism of graphs, we may choose  $k < l$  or  $k = l = 1$ .

Since  $\Gamma$  is connected,  $G = \langle S_k \rangle = \langle c \rangle \times \langle ab^k, b^l \rangle$ , we have  $\langle ab^k, b^l \rangle = \langle a, b \rangle$ . Since  $\langle a \rangle$  is normal in  $\langle a, b \rangle$ , we may set  $a^{b^l} = a^e$  for some integer  $e$ . Since  $o(a) = m$ , we have  $e^2 \equiv 1 \pmod{m}$ , and so  $H := \langle a, b \rangle = \langle ab^k, b^l \rangle = \langle a^e b^k, ab^k, b^l \rangle = \langle a^{e-1}, ab^k, b^l \rangle = \langle a^{e-1} \rangle \langle ab^k, b^l \rangle$ . Let  $K = \langle a^{e-1} \rangle$ . Since  $(ab^k)^{b^l} = a^e b^k = a^{e-1} ab^k$ , we have  $K(ab^k)^{b^l} = Ka^{e-1} ab^k = Kab^k$ . Thus, the quotient group  $H/K$  is abelian, so  $\langle a \rangle = H' \leq K = \langle a^{e-1} \rangle$ . Then  $\langle a \rangle = \langle a^{e-1} \rangle$ , and so  $(e-1, m) = 1$ . Hence,  $e \equiv -1 \pmod{m}$  as  $e^2 \equiv 1 \pmod{m}$ , and so  $a^{b^l} = a^{-1}$ .

Now we show that  $\text{Aut}(G, S_k) \neq 1$  if and only if  $l = 1$ . Suppose that  $\text{Aut}(G, S_k) \neq 1$ . Then, since  $S_k$  contains only one involution, we conclude that  $\text{Aut}(G, S_k) = \langle \tau \rangle \cong \mathbb{Z}_2$ ,  $b^l = (b^l)^\tau$  and  $(cab^k)^\tau = (cab^k)^{-1}$ . Then  $c^\tau = c^{-1}$  and  $(ab^k)^\tau = (ab^k)^{-1} = b^{-k} a^{-1} = (a^{-1})^{b^k} b^{-k} = a^s b^{-k}$  for some  $s$ . By [14], we set  $a^\tau = a^i$  and  $b^\tau = a^j b$  for some  $i$  and  $j$ . Then, noting  $a^b \in \langle a \rangle$ , computation shows that  $(ab^k)^\tau = a^\tau (b^\tau)^k = a^{i+tk} b^k$  for some  $t$ . Thus  $a^{i+tk} b^k = a^s b^{-k}$ , yielding  $k \equiv -k \pmod{2l}$ , and so  $l = 1$  as  $(l, k) = 1$ .

Conversely, suppose that  $l = 1$ . Then  $o(c) = \frac{2n}{o(a)o(b)} = \frac{n}{m} > 1$ ,  $k = 0$  or  $1$ , and  $S_k = \{ca, c^{-1}a^{-1}, b\}$  or  $\{cab, c^{-1}ab, b\}$ . Assume first that  $S_k = \{cab, c^{-1}ab, b\}$ . Take  $\tau \in \text{Aut}(G)$  with  $c^\tau = c^{-1}$ ,  $a^\tau = a$ , and  $b^\tau = b$ . Then  $1 \neq \tau \in \text{Aut}(G, S_k)$ , and  $\text{AutCay}(G, S_k)$  has a regular subgroup  $\langle \bar{c}\bar{a}, \bar{b}\tau \rangle \cong D_{2n}$ , so  $\Gamma$  is a dihedrant. Now let  $S_k = \{ca, c^{-1}a^{-1}, b\}$ . By [14], take  $\tau \in \text{Aut}(G)$  with  $c^\tau = c^{-1}$ ,  $a^\tau = a^{-1}$ , and  $b^\tau = b$ . Then  $1 \neq \tau \in \text{Aut}(G, S_k)$ . Since  $\langle ca \rangle$  is normal in  $G$ , we set  $(ca)^b = (ca)^t$  for some  $1 < t < n$ . Then  $t^2 \equiv 1 \pmod{n}$  as  $o(b) = 2$  and  $o(ca) = n$ . Let  $r = t$  or  $n - t$  such that  $r < \frac{n}{2}$ . For  $0 \leq i \leq n - 1$ , we label  $\alpha_i = (ca)^i$  and  $\beta_i = b(ca)^i$  if  $r = t$ , or  $\alpha_i = (ca)^{-i}$  and  $\beta_i = b(ca)^{-i}$  if  $r = n - t$ . Then  $\text{Cay}(G, S_k)$  has edges  $\{\alpha_i, \alpha_{i+1}\}$ ,  $\{\alpha_i, \beta_i\}$ , and  $\{\beta_i, \beta_{i+r}\}$ . Thus  $\Gamma \cong \text{Cay}(G, S_k) \cong \mathbf{P}(n, r)$ . ■

### 3. CUBIC COSET GRAPHS

In a graph, an *arc* is an ordered pair of adjacent vertices, and a *2-arc* is a directed path of length 2. A graph  $\Gamma$  is called *arc-transitive* or *2-arc-transitive* if  $\text{Aut}\Gamma$  is transitive on the arcs or the 2-arcs of  $\Gamma$ , respectively. For a graph  $\Gamma$  and  $G \leq \text{Aut}\Gamma$ , we say  $\Gamma$  to be *G-vertex-transitive* or *G-arc-transitive* if  $G$  acts transitively on the vertices or the arcs of  $\Gamma$ , respectively.

Let  $\Gamma = (V, E)$  be a  $G$ -vertex-transitive graph. Then, for  $\alpha \in V$ , the stabilizer  $G_\alpha$  is a core-free subgroup in  $G$ , that is,  $\bigcap_{g \in G} G_\alpha^g = 1$ . Set  $H = G_\alpha$  and  $D = \{x \mid \alpha^x \in \Gamma(\alpha)\}$ , where  $\Gamma(\alpha)$  is the set of neighbors of  $\alpha$  in  $\Gamma$ . Then  $D$  is a union of several double cosets  $HxH$ . Since  $\Gamma$  is undirected, we have  $D = D^{-1} := \{x^{-1} \mid x \in D\}$ . Moreover,  $\Gamma$  is isomorphic to the *coset graph*  $\text{Cos}(G, H, D)$  defined over  $\{Hx \mid x \in G\}$  with edge set  $\{\{Hg_1, Hg_2\} \mid g_2g_1^{-1} \in D\}$ .

The following statements for coset graphs are well known.

- (a)  $\Gamma$  is connected if and only if  $\langle H, D \rangle = G$ .
- (b)  $\Gamma$  is  $G$ -arc-transitive if and only if  $D = HgH$  for  $g \in G$  with  $g^2 \in H$ ; moreover,  $g$  can be chosen as a 2-element with  $g \in \mathbf{N}_G(H \cap H^g)$  and  $g^2 \in H \cap H^g$ .

The next lemma gives a characterization of the prime divisors of  $|G_\alpha|$ .

**Lemma 3.1** ([7]). *If  $\Gamma$  is connected and of valency  $k$ , then each prime divisor of  $|G_{\alpha\beta}|$  is less than  $k$ , where  $\{\alpha, \beta\}$  is an edge of  $\Gamma$ .*

Now assume that  $\Gamma$  is cubic and connected. If  $G$  is regular on  $V$ , then  $\Gamma$  is a Cayley graph of  $G$ . If  $G$  is transitive on the arcs of  $\Gamma$ , then  $\Gamma \cong \text{Cos}(G, G_\alpha, G_\alpha g G_\alpha)$  where  $g$  is a 2-element with  $\langle g, G_\alpha \rangle = G$ ,  $\alpha^g \in \Gamma(\alpha)$ ,  $g \in \mathbf{N}_G(G_{\alpha\alpha^g})$ , and  $g^2 \in G_{\alpha\alpha^g}$ ; moreover, the well-known result of Tutte determines  $G_\alpha$ , refer to [2].

**Theorem 3.2.** *If  $\Gamma$  is  $G$ -arc-transitive, then  $G_\alpha \cong \mathbb{Z}_3, S_3, D_{12}, S_4$  or  $S_4 \times S_2$ .*

Suppose that  $G$  is not regular on  $V$  and not transitive on the arcs of  $\Gamma$ . Then  $G_\alpha$  fixes one of neighbors, say  $\gamma$ , and transitive on the other two neighbors, say  $\beta_1$  and  $\beta_2$ , of  $\alpha$ . Thus  $G_\alpha$  is a nontrivial 2-group by Lemma 3.1. Moreover,  $\Gamma$  is an arc-disjoint union of two  $G$ -arc-transitive graphs, one of valency 2 and the other of valency 1. Then  $\Gamma \cong \text{Cos}(G, G_\alpha\{x, y\}G_\alpha)$ , where  $x$  and  $y$  are 2-elements such that  $\alpha = \beta_1^x$ ,  $x \in \mathbf{N}_G(G_{\alpha\beta_1})$ ,  $x^2 \in G_{\alpha\beta_1}$ ,  $\alpha^y = \gamma$ ,  $y \in \mathbf{N}_G(G_\alpha)$ ,  $y^2 \in G_\alpha$ , and  $\langle x, y, G_\alpha \rangle = G$ . Thus, if a characteristic subgroup  $M \leq G_{\alpha\beta_1}$  is normal in  $\langle y, G_\alpha \rangle$  then  $M = 1$ ; if  $G$  has an abelian Sylow

2-subgroup, then  $\langle y, G_\alpha \rangle$  is an abelian 2-group, and so  $G_{\alpha\beta_1}$  is normal in  $G$ , hence  $G_{\alpha\beta_1} = 1$ . Then the next lemma follows.

**Lemma 3.3.** *Assume that  $\{\beta_1, \beta_2\}$  and  $\{\gamma\}$  are the two  $G_\alpha$ -orbits on  $\Gamma(\alpha)$ . Then  $G_\alpha$  and  $G_{\alpha\beta_1}$  do not contain a common nontrivial characteristic subgroup. If further  $G$  has an abelian Sylow 2-subgroup, then  $G_\alpha \cong \mathbb{Z}_2$ .*

Some of the generalized Petersen graphs can be constructed as coset graphs.

**Lemma 3.4.** *Let  $\Gamma$  be a connected  $G$ -vertex-transitive cubic graph with  $\mathbb{Z}_n : \mathbb{Z}_4 \cong G \leq \text{Aut}\Gamma$ , where  $n$  is odd and square-free. Then either  $G$  is a regular subgroup of  $\text{Aut}\Gamma$ , or  $\Gamma \cong \mathbf{P}(n, r)$  for  $1 < r < \frac{n}{2}$  with  $r^2 \equiv -1 \pmod{n}$ .*

**Proof.** Let  $\langle a \rangle$  be the normal subgroup of  $G$  of order  $n$ . Then  $\langle a \rangle$  is a semiregular subgroup of  $G$ . Since  $\langle a \rangle$  has odd order and  $\Gamma$  has valency 3, we conclude that  $\langle a \rangle$  is intransitive on  $V\Gamma$ . Thus,  $\Gamma$  has order  $2n$  or  $4n$ . If  $\Gamma$  has order  $4n$ , then  $G$  is a regular subgroup of  $\text{Aut}\Gamma$ . Hence, we assume  $\Gamma$  has order  $2n$ . Let  $b \in G$  be of order 4. Then  $G = \langle a \rangle : \langle b \rangle$  and  $a^b = a^r$  as  $\langle a \rangle$  normal in  $G$ , where  $1 \leq r < n$  with  $r^4 \equiv 1 \pmod{n}$ .

Note all involutions of  $G$  are conjugate and contained in  $\langle a, b^2 \rangle$ . Then  $H := G_\alpha = \langle b^2 \rangle$  for some  $\alpha \in V\Gamma$ . Write  $\Gamma \cong \text{Cos}(G, H, H\{x, y\}H)$ , where  $x$  is an involution and  $y \in \mathbf{N}_G(H)$  with  $y^2 \in H$ . Let  $\mathbf{C}_{\langle a \rangle}(b^2) = \langle a_1 \rangle$ . Since  $o(a) = n$  is square-free, we may write  $\langle a \rangle = \langle a_1 \rangle \times \langle a_2 \rangle$ . Then  $a_2 \neq 1$ ; otherwise,  $\mathbf{C}_{\langle a \rangle}(b^2) = \langle a \rangle$ , yielding  $H = \langle b^2 \rangle$  is normal in  $G$ , a contradiction. It is easily shown that  $a_2^{b^2} = a_2^{-1}$ , yielding  $a_2^2 = a_2^{-1}$ , and hence  $r^2 \equiv -1 \pmod{o(a_2)}$ . Note that  $\mathbf{N}_G(H) = \langle a_1 \rangle : \langle b \rangle$  and all involutions of  $G$  are contained in  $\langle a_2, b^2 \rangle$ . Since  $HbH = Hb^{-1}H$  and  $\langle x, y, H \rangle = G$ , we may choose  $x = a_2^t b^2$  and  $y = a_1^i b$  with  $y^2 \in H$ . Then  $y^2 = a_1^i b^2 (b^{-1} a_1^i b) = a_1^{i+ri} b^2$ , yielding  $y^2 = b^2$ . In particular,  $y$  has order 4. Thus, since  $\Gamma$  is connected,  $G = \langle x, y, H \rangle = \langle a_2^t b^2, y, y^2 \rangle = \langle a_2^t, y \rangle = \langle a_2^t \rangle : \langle y \rangle$ . It follows that  $\langle a \rangle = \langle a_2^t \rangle$ , and so  $n = o(a) = o(a_2) = o(a_2^t)$ ,  $a_1 = 1$ , and  $r^2 \equiv -1 \pmod{n}$ . Thus  $y = b$ , and it is easily shown that  $\mathbf{N}_G(H) = \langle b \rangle$ . Write  $a_2^t = a^s$ . Then  $x = a^s b^2$  and  $G = \langle a^s \rangle : \langle b \rangle$ .

Since  $H\{x, y\}H = H\{a^s, b\}H$ , we have  $\Gamma \cong \text{Cos}(G, H, H\{a^s, b\}H)$ . Since  $HbH = Hb^3H$  and  $a^{b^3} = a^{n-r}$ , replacing  $b$  by  $b^3$  if necessary, we assume that  $r < \frac{n}{2}$ .

Now label  $\alpha_i = Ha^{si}$  and  $\beta_i = Hba^{si}$ , where  $0 \leq i \leq n-1$ , which gives rise to all vertices of  $\Gamma$ . Then,  $\{\alpha_i, \alpha_{i+1}\}$  and  $\{\alpha_i, \beta_i\}$  are edges. Moreover,  $\beta_i = Hba^{si}$  and  $\beta_j = Hba^{sj}$  are adjacent whenever  $(a^s)^{(j-i)(-r)} = ba^{sj-si}b^{-1} = ba^{sj}(ba^{si})^{-1}$  equals to  $a^s$  or  $a^{-s}$ , i.e.,  $(j-i)(-r) \equiv \pm 1 \pmod{n}$ . Thus  $\{\beta_i, \beta_j\}$  is an edge if and only if  $j \equiv i \pm r \pmod{n}$ . Therefore,  $\Gamma \cong \text{Cos}(G, H, H\{a^s, b\}H) \cong \mathbf{P}(n, r)$ . ■

We next describe some graphs associated with simple groups  $\text{PSL}(2, p)$  with  $p$  prime. As usual, for two integers  $d, n$ , by  $d \parallel n$  we mean  $d$  divides  $n$ , and  $(d, \frac{n}{d}) = 1$ .

**Example 3.5.** Let  $T = \text{PSL}(2, p)$ , where  $p$  is a prime.

- (1) Assume that  $p \equiv \pm 3 \pmod{8}$ . Then  $4 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Take a subgroup  $H \cong S_3$  of  $T$ , and let  $K \cong \mathbb{Z}_2$  be a Sylow 2-subgroup of  $H$ . Then  $\mathbf{N}_T(K) = D_{p-\varepsilon}$ , and let  $g \in \mathbf{N}_T(K) \setminus K$  be an involution such that  $\langle H, g \rangle = T$ .
- (2) Assume that  $p \equiv \pm 7 \pmod{16}$ . Then  $8 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Take a subgroup  $H \cong D_{12}$  of  $T$ , and let  $K \cong \mathbb{Z}_2^2$  be a Sylow 2-subgroup of  $H$ . Then  $\mathbf{N}_T(K) = S_4$ , and let  $g \in \mathbf{N}_T(K) \setminus K$  be an involution such that  $\langle H, g \rangle = T$ .

- (3) Assume that  $p \equiv \pm 15 \pmod{32}$ . Then  $16 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Take a subgroup  $H \cong S_4$  of  $T$ , and let  $K \cong D_8$  be a Sylow 2-subgroup of  $H$ . Then  $N_T(K) = D_{16}$ , and let  $g \in N_T(K) \setminus K$  be an involution such that  $\langle H, g \rangle = T$ .

In each of these three cases, the coset graph  $\Gamma = \text{Cos}(T, H, HgH)$  is a connected 2-arc-transitive cubic graph, and the order of  $\Gamma$  is even and indivisible by 4.

**Example 3.6.** Let  $T = \text{PSL}(2, p)$ , and let  $G = \text{PGL}(2, p)$ , where  $p$  is a prime.

- (1) Assume that  $p \equiv \pm 3 \pmod{8}$ . Then  $4 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Take a subgroup  $H \cong D_{12}$  of  $T$ , and let  $K \cong \mathbb{Z}_2^2$  be a Sylow 2-subgroup of  $H$ . Then  $N_G(K) = S_4$ . Let  $g \in N_G(K) \setminus K$  be an involution such that  $\langle H, g \rangle = G$ .
- (2) Assume that  $p \equiv \pm 7 \pmod{16}$ . Then  $8 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Take a subgroup  $H \cong S_4$  of  $T$ , and let  $K \cong D_8$  be a Sylow 2-subgroup of  $H$ . Then  $N_G(K) = D_{16}$ , and let  $g \in N_G(K) \setminus K$  be an involution such that  $\langle H, g \rangle = G$ .

If  $g$  is described as in (1) or (2), then the coset graph  $\Gamma = \text{Cos}(G, H, HgH)$  is bipartite, connected, cubic, and 2-arc-transitive.

The final two examples give several families of cubic graphs associated with  $\text{PSL}(2, p)$ , which are not arc-transitive.

**Example 3.7.** Let  $T = \text{PSL}(2, p)$ , where  $p$  is a prime.

- (1) Assume that  $p \equiv \pm 3 \pmod{8}$ . Then  $4 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Let  $\mathbb{Z}_2 \cong H < T$ . Then  $N_T(H) = D_{p-\varepsilon}$ . Let  $x \in N_T(H) \setminus H$  and  $y \in T \setminus N_T(H)$  be two involutions. Then  $\langle H, x, y \rangle = T$ .
- (2) Assume that  $p \equiv \pm 7 \pmod{16}$ . Then  $8 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Let  $\mathbb{Z}_2^2 \cong H < T$ , and let  $K \cong \mathbb{Z}_2$  be a subgroup of  $H$ . Then  $N_T(H) = S_4$  and  $N_T(K) = D_{p-\varepsilon}$ . Let  $x \in N_T(H) \setminus H$  and  $y \in N_T(K) \setminus N_{N_T(H)}(K)$  be involutions such that  $\langle H, x, y \rangle = T$ .
- (3) Assume that  $p \equiv \pm 15 \pmod{32}$ . Then  $16 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Let  $D_8 \cong H < T$  and  $K \cong \mathbb{Z}_2^2$  be a subgroup of  $H$ . Then  $N_T(H) = D_{16}$  and  $N_T(K) = S_4$ . Let  $x \in N_T(H) \setminus H$  and  $y \in N_T(K) \setminus H$  be involutions such that  $\langle H, x, y \rangle = T$ .

Take  $x$  and  $y$  as in (1), (2), or (3). Then the coset graph  $\Gamma = \text{Cos}(T, H, H\{x, y\}H)$  is a connected cubic graph, and  $\Gamma$  has even order indivisible by 4.

**Example 3.8.** Let  $T = \text{PSL}(2, p)$ , and let  $G = \text{PGL}(2, p)$ , where  $p$  is a prime.

- (1) Assume that  $p \equiv \pm 3 \pmod{8}$ . Then  $4 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Let  $\mathbb{Z}_2^2 \cong H < T$  and  $K \cong \mathbb{Z}_2$  be a subgroup of  $H$ . Then  $N_G(K) = D_{2((p-\varepsilon))}$  and  $N_G(H) = S_4$ . Let  $x \in N_G(H) \setminus H$  and  $y \in N_G(K) \setminus N_{N_G(K)}(H)$  be two involutions such that  $\langle H, x, y \rangle = G$ .
- (2) Assume that  $p \equiv \pm 7 \pmod{16}$ . Then  $8 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Let  $D_8 \cong H < T$  and let  $K \cong \mathbb{Z}_2^2$  be a subgroup of  $H$ . Then  $N_G(H) = D_{16}$  and  $T > N_G(K) = S_4$ . Let  $x \in N_G(K) \setminus H$  and  $y \in N_G(K) \setminus H$  be an involution such that  $\langle H, x, y \rangle = G$ .

For each of (1) and (2), the coset graph  $\Gamma = \text{Cos}(G, H, H\{x, y\}H)$  is bipartite, connected, and cubic, and the order of  $\Gamma$  is even and indivisible by 4.

#### 4. NORMAL QUOTIENTS

Let  $\Gamma = (V, E)$  be a connected  $G$ -vertex-transitive graph, where  $G \leq \text{Aut}\Gamma$ .

For a normal subgroup  $N \triangleleft G$ , the *normal quotient*  $\Gamma_N$  of  $\Gamma$ , induced by  $N$ , is the graph whose vertices are the  $N$ -orbits on  $V$  such that  $B$  and  $C$  are adjacent if and only if there exists an edge  $\{\beta, \gamma\} \in E$  with  $\beta \in B$  and  $\gamma \in C$ . Clearly, the valency of  $\Gamma_N$  is at most the number of  $N_\alpha$ -orbits on  $\Gamma(\alpha)$ . Let  $K$  be the kernel of  $G$  acting on the  $N$ -orbits. Then  $G/K$  can be viewed as a subgroup of  $\text{Aut}\Gamma_N$ . If the valency of  $\Gamma_N$  equals the valency of  $\Gamma$ , then  $\Gamma$  is a *cover* of  $\Gamma_N$  and, in this case,  $K = N$  is semiregular on  $V$ .

From now on, we assume that  $\Gamma$  is connected and cubic. Suppose that  $G$  is neither regular on  $V$  nor transitive on the arcs of  $\Gamma$ . Then  $G_\alpha$  is a nontrivial 2-group, where  $\alpha \in V$ . Set  $\Gamma(\alpha) = \{\beta_1, \beta_2, \gamma\}$  such that  $G_\alpha$  is transitive on  $\{\beta_1, \beta_2\}$  and fixes  $\gamma$ .

Let  $N \triangleleft G$  have at least three orbits on  $V$ , and  $V_N$  be the set of  $N$ -orbits. Then the quotient graph  $\Gamma_N$  has valency 2 or 3. If  $\Gamma_N$  has valency 3, then  $\Gamma$  is a cover of  $\Gamma_N$ .

**Lemma 4.1.** *Let  $K$  be the kernel of  $G$  acting on  $V_N$ . If  $\Gamma$  is not a cover of  $\Gamma_N$ , then  $\Gamma_N$  is an  $l$ -cycle and either*

- (1) *each  $N$ -orbit is a matching,  $K = N$  is semiregular,  $G/N \cong D_{2l}$ , and  $G$  has a regular subgroup  $N.\mathbb{Z}_l$ ; or*
- (2)  *$G_\alpha = K_\alpha$  is a 2-group,  $l$  is even, and  $G/K \cong D_l$  acting on  $V_N$  regularly.*

**Proof.** Suppose that  $\Gamma_N$  has valency 2. Then  $\Gamma_N$  is an  $l$ -cycle for some integer  $l$ . Noting that  $(\gamma^N)^{G_\alpha} = \gamma^N$  and  $(\beta_1^N)^g = \beta_2^N$  for some  $g \in G_\alpha$ , either  $\alpha^N = \gamma^N$  and  $\beta_1^N \neq \beta_2^N$ , or  $\alpha^N \neq \gamma^N$  and  $\beta_1^N = \beta_2^N$ .

We assume first that  $\alpha^N = \gamma^N$  and  $\beta_1^N \neq \beta_2^N$ . Then  $\alpha^N$  induces a matching, and  $G/K$  is transitive on the arcs of  $\Gamma_N$ , and so  $G/K \cong D_{2l}$ . Noting that  $K_\alpha$  fixes  $\Gamma(\alpha) = \{\beta_1, \beta_2, \gamma\}$  point-wise, it implies that  $K_\alpha = 1$ , hence  $N = K$  is a semiregular subgroup of  $G$ . Then  $G$  contains a subgroup  $N.\mathbb{Z}_l$  which is regular on  $V$ .

Now let  $\alpha^N \neq \gamma^N$  and  $\beta_1^N = \beta_2^N$ . Then the induced subgraphs  $[\alpha^N \cup \beta_1^N]$  and  $[\alpha^N \cup \gamma^N]$  are regular and have valency 2 and 1, respectively. Thus, there is no an element in  $G$  which maps  $\{\alpha^N, \beta_1^N\}$  to  $\{\alpha^N, \gamma^N\}$ . Therefore,  $G/K$  is transitive on  $V_N$  but not on the edges of  $\Gamma_N$ . Noting that  $\text{Aut}\Gamma_N \cong D_{2l}$ , it follows that  $l$  is even,  $G/K \cong D_l$  and  $G/K$  acting on  $V_N$  regularly. Moreover,  $K_\alpha = G_\alpha$ . ■

This leads us to define a special type of cover for some cubic graphs.

**Construction 4.2.** Assume that  $X = \text{PGL}(2, p)$ ,  $T = \text{PSL}(2, p)$ , and  $p \equiv \pm 3 \pmod{8}$ . Then  $4 \parallel (p - \varepsilon)$ , where  $\varepsilon = 1$  or  $-1$ . Let  $\mathbb{Z}_2^2 \cong H < T$  and  $K \cong \mathbb{Z}_2$  be a subgroup of  $H$ . Then  $N_X(K) = D_{2((p-\varepsilon))}$  and  $N_X(H) = S_4$ . Let  $x \in N_X(H) \setminus T$  and  $y \in N_X(K) \setminus T$  be such that  $x^2 \in H$ ,  $y^2 \in K$ , and  $\langle H, x, y \rangle = X$ . Let  $M = \langle c \rangle \cong \mathbb{Z}_m$  with odd  $m$  coprime to  $|T|$ , and let  $G = (T \times M)\langle x \rangle$  such that  $c^x = c^{-1}$  (and so  $c^y = c^{-1}$ ). Then  $G = T:D_{2m}$ , and  $\Sigma = \text{Cos}(G, H, H\{c^i x, c^j y\}H)$  is a cubic graph.

It is easily shown that  $\Sigma$  is connected if and only if  $(i - j, m) = 1$ . Moreover,  $\Sigma_M \cong \text{Cos}(X, H, H\{x, y\}H)$  and  $\Sigma_T$  is a cycle of length  $2m$ .

## 5. SOLUBLE AUTOMORPHISM GROUPS

Let  $\Gamma = (V, E)$  be a connected cubic  $G$ -vertex-transitive graph of square-free order  $2n$ , where  $G \leq \text{Aut}\Gamma$ . In this section, we consider the case where  $G$  is soluble.

If  $G$  is regular on  $V$ , then  $\Gamma$  is a Cayley graph of  $G$ , and  $\Gamma$  is known by Lemmas 2.1–2.5 and Corollary 2.4. Thus, in the following, we assume that  $G$  is not regular on  $V$ , that is,  $G_\alpha \neq 1$  for  $\alpha \in V$ . Then Lemma 4.1 is available.

As usual, for a prime divisor  $p$  of  $|G|$ , let  $\mathbf{O}_p(G)$  be the largest normal  $p$ -subgroup of  $G$ . Since the order  $|G : G_\alpha|$  of  $\Gamma$  is square-free and  $G_\alpha$  is a  $\{2, 3\}$ -group, either  $|\mathbf{O}_p(G)| \leq p$ , or  $|\mathbf{O}_p(G)| \geq p^2$  and  $p \in \{2, 3\}$ .

**Lemma 5.1.** *If  $\mathbf{O}_2(G) \neq 1$ , then  $G \cong \mathbb{Z}_{2n}:\mathbb{Z}_2 \cong D_{4n}$ , and  $\Gamma = \mathbf{M}_n$  or  $\mathbf{P}(n, 1)$ .*

**Proof.** Let  $N = \mathbf{O}_2(G) \neq 1$ . Then each  $N$ -orbit has length 2, and the quotient graph  $\Gamma_N$  is of odd order  $n$ . It follows from Lemma 4.1 that  $G_\alpha \cong \mathbb{Z}_2$ ,  $N \cong \mathbb{Z}_2$  and  $G$  contains a regular subgroup  $N.\mathbb{Z}_n \cong \mathbb{Z}_{2n}$ , and so  $G \cong \mathbb{Z}_{2n}:\mathbb{Z}_2$ . Thus  $G$  contains a normal regular subgroup  $R \cong \mathbb{Z}_{2n}$ . Write  $\Gamma = \text{Cay}(R, S)$ . Then  $S = \{a, a^{-1}, b\}$ , where  $b$  is the unique involution in  $R$ , and  $o(a) = n$  or  $2n$ . Thus,  $\Gamma = \mathbf{M}_n$  or  $\mathbf{P}(n, 1)$ .

Let  $\alpha$  be the vertex corresponding the identity of  $R$ . Then  $G_\alpha \leq \text{Aut}(R)$ . Set  $G_\alpha = \langle \sigma \rangle$ . Then  $a^\sigma = a^{-1}$  as  $S^\sigma = S$ , and thus  $G = R:\langle \sigma \rangle \cong D_{4n}$ . ■

**Lemma 5.2.** *If  $\mathbf{O}_3(G)$  has order divisible by 9, then  $\Gamma = \mathbf{K}_{3,3}$  and  $\text{Aut}\Gamma = \mathbf{S}_3 \wr \mathbf{S}_2$ .*

**Proof.** Let  $N = \mathbf{O}_3(G)$ . Assume that  $|N| > 3$ . Then  $N$  is not semiregular on  $V$ , and  $N_\alpha$  is a nontrivial 3-group. It follows that  $N_\alpha$  is transitive on  $\Gamma(\alpha)$ . For  $\beta \in \Gamma(\alpha)$ , the orbit  $\beta^{N_\alpha}$  has size 3. It follows that the induced subgraph of  $\Gamma$  with vertex set  $\alpha^N \cup \beta^N$  is isomorphic to  $\mathbf{K}_{3,3}$ . So  $\Gamma \cong \mathbf{K}_{3,3}$ , and clearly,  $\text{Aut}\Gamma = \mathbf{S}_3 \wr \mathbf{S}_2$ . ■

Let  $F$  be the Fitting subgroup of  $G$ , the largest nilpotent normal subgroup of  $G$ . Then  $F \neq 1$  and  $\mathbf{C}_G(F) \leq F$  as  $G$  is soluble, and  $F = \langle \mathbf{O}_p(G) \mid p \mid |G| \rangle$ .

**Lemma 5.3.** *Assume that  $\mathbf{O}_2(G) = 1$  and  $\mathbf{O}_3(G) = 1$  or  $\mathbb{Z}_3$ . Then Fitting subgroup of  $G$  is cyclic and has exactly two orbits on  $V$ , and either  $\Gamma \cong \mathbf{K}_{3,3}$  or one of the following holds.*

- (1)  $\mathbb{Z}_n:\mathbb{Z}_4$  and  $\Gamma \cong \mathbf{P}(n, r)$ , where  $r^2 \equiv -1 \pmod{n}$ ;
- (2)  $G \cong \mathbb{Z}_n:\mathbb{Z}_2^2$  and  $\Gamma \cong \mathbf{M}_n$  or  $\mathbf{P}(n, r)$ , where  $r^2 \equiv 1 \pmod{n}$ ;
- (3)  $G \cong \mathbb{Z}_n:\mathbb{Z}_6 \cong D_{2n}:\mathbb{Z}_3$  and  $\Gamma$  is isomorphic to one of the graphs involved in Lemma 2.3 (3).

**Proof.** Let  $F$  be the Fitting subgroup of  $G$ . Noting that  $\mathbf{O}_2(G) = 1$  and  $\mathbf{O}_p(G) = 1$  or  $\mathbb{Z}_p$  for each odd prime  $p$  divisor of  $|G|$ , we conclude that  $F$  is cyclic and of odd order. It follows that  $F$  is semiregular on  $V$ . Since  $\mathbf{C}_G(F) \leq F$ , we have  $\mathbf{C}_G(F) = F$ . Then  $G/F = \mathbf{N}_G(F)/\mathbf{C}_G(F)$  is isomorphic to a subgroup of  $\text{Aut}(F)$ , which is abelian.

Suppose that  $F$  has at least three orbits on  $V$ . Then, by Lemma 4.1,  $\Gamma$  is a cover of  $\Gamma_F$ . Thus  $G/F$  is isomorphic to a subgroup of  $\text{Aut}\Gamma_F$ , and so  $G/F$  is regular on  $V_F$  as it is abelian. Then  $G$  is regular on  $V$ , which is not the case.

Thus,  $F$  has at most two orbits on  $V$ . Since  $F$  has odd order,  $F$  has exactly two orbits on  $V$ . Since  $G/F$  is abelian,  $G$  has an abelian Sylow 2-subgroup. If  $G$  is not transitive on the arcs of  $\Gamma$ , then  $G_\alpha \cong \mathbb{Z}_2$  by Lemma 3.3, and so  $G = F:\mathbb{Z}_2^2$  or  $F:\mathbb{Z}_4$ . On the other

hand,  $G_\alpha \cong G_\alpha/F_\alpha \cong FG_\alpha/F \leq G/F$  is abelian. If  $\Gamma$  is  $G$ -arc-transitive, then  $G_\alpha \cong \mathbb{Z}_3$  by Theorem 3.2, so  $G = F:\mathbb{Z}_6$ . If  $G \cong \mathbb{Z}_n:\mathbb{Z}_4$  then (1) holds by Lemma 3.4. If  $G \not\cong \mathbb{Z}_n:\mathbb{Z}_4$  then  $G$  has a normal regular subgroup  $R \cong \mathbb{Z}_n:\mathbb{Z}_2$ , and so  $\Gamma$  is known by Lemmas 2.1–2.5 and Corollary 2.4. This completes the proof. ■

## 6. INSOLUBLE AUTOMORPHISM GROUPS

Let  $\Gamma = (V, E)$  be a connected cubic  $G$ -vertex-transitive graph of square-free order  $2n$ , where  $G \leq \text{Aut}\Gamma$ . In this section, we assume that  $G$  is insoluble.

Recall that the *soluble radical* of a group  $G$  is the largest soluble normal subgroup of  $G$ . Since  $G$  is insoluble, the next lemma is a consequence of Lemma 4.1.

**Lemma 6.1.** *Let  $M$  be the soluble radical of  $G$ . Then  $\Gamma$  is a cover of  $\Gamma_M$ ; in particular,  $M$  is semiregular on  $V$  and of odd order.*

**Proof.** Let  $V_M$  be the set of  $M$ -orbits on  $V$ , and let  $K$  be the kernel of  $G$  acting on  $V_M$ . Then  $M \triangleleft K \triangleleft G$ , and  $K = MK_\alpha$ . Since  $K_\alpha \triangleleft G_\alpha$  is soluble, so is  $K$ , and hence  $K = M$ . Thus,  $G/M \leq \text{Aut}\Gamma_M$  is insoluble, and so  $\Gamma_M$  is cubic. Hence  $M$  is semiregular, and  $|V_M|$  is even. Since  $|V| = |M||V_M|$  is square-free,  $|M|$  is odd. ■

We first deal with the case where  $G$  has trivial soluble radical.

**Lemma 6.2.** *Suppose that the soluble radical of  $G$  is trivial. Then  $G$  is almost simple.*

**Proof.** Let  $N$  be a minimal normal subgroup of  $G$ . Then  $N$  is insoluble. Let  $V_N$  be the set of  $N$ -orbits on  $V$ , and let  $K$  be the kernel of  $G$  on  $V_N$ . Then  $K = NK_\alpha$ , and so  $K/N$  is soluble. Since  $|V|$  is square-free,  $N$  is not semiregular on  $V$ , and hence the quotient graph  $\Gamma_N$  has valency 0, 1, or 2. Thus,  $G/K \leq \text{Aut}\Gamma_N$  is soluble, and so is  $G/N$ . Hence  $N$  is the only minimal normal subgroup of  $G$ . Since  $|G|$  is not divisible by  $p^2$  with  $p \geq 5$  prime,  $N$  is simple, and  $G$  is almost simple. ■

**Lemma 6.3.** *Let  $G$  be almost simple with socle  $\text{soc}(G) = T$ . Assume that  $\Gamma$  is  $G$ -arc-transitive. Then either*

- (1)  $T = A_6$ ,  $\text{Aut}\Gamma = \text{P}\Gamma\text{L}(2, 9)$  and  $\Gamma$  is isomorphic to Tutte's 8-cage, or
- (2)  $T = \text{PSL}(2, p)$  such that a Sylow 2-subgroup of  $T$  is  $\mathbb{Z}_2^2$ ,  $D_8$ , or  $D_{16}$ , and  $\Gamma$  is a 2-arc-transitive graph; moreover,  $\Gamma$  is described as in Example 3.5 or 3.6.

**Proof.** By Theorem 3.2,  $|G_\alpha|$  is not divisible by  $2^5 \cdot 3^2$ . Since  $|V| = |G : G_\alpha|$  is square-free,  $|G|$  is not divisible by  $2^6$ ,  $3^3$ , and  $r^2$ , where  $r$  is a prime with  $r > 3$ . Inspecting the orders of finite simple groups, we obtain that  $T$  is one of  $A_6$ ,  $A_7$ ,  $M_{11}$ ,  $J_1$ ,  $\text{PSL}(2, 2^f)$ ,  $\text{PSL}(2, p)$  for prime  $p \geq 5$ .

Suppose that  $T = \text{PSL}(2, 2^f)$  with  $f \geq 3$ . Then  $f = 3, 4$ , or  $5$ . By the information given in the Atlas [8], we conclude that  $G$  has no a subgroup of square-free index as listed in Theorem 3.2, which is a contradiction.

Suppose that  $T = A_7$ . Note that  $|G : G_\alpha|$  is even and square-free. Then either  $|T_\alpha| = 12$  and  $T$  is transitive on  $V$ , or  $|G_\alpha| = |T_\alpha| = 24$  and  $T$  has two orbits on  $V$ . Thus,  $\Gamma$  is a  $G$ -arc-transitive graph of order 210; however, by [6], there exists no such a graph, which is a contradiction.

Suppose that  $T = M_{11}$ . Then  $G = T$  and  $|T_\alpha| = 24$ , so  $T_\alpha \cong S_4$ . Thus,  $T_{\alpha\beta} \cong D_8$  and  $N_T(T_{\alpha\beta})$  is a Sylow 2-subgroup of  $T$ , where  $\beta \in \Gamma(\alpha)$ . Further, computation using GAP shows that all subgroups of  $T$  isomorphic to  $S_4$  are conjugate. Thus we may assume that  $T_\alpha$  is contained in a maximal subgroup  $M \cong M_{10}$ . So  $N_T(T_{\alpha\beta}) = N_M(T_{\alpha\beta})$ . Then there is no  $x \in N_T(T_{\alpha\beta})$  with  $\langle x, T_\alpha \rangle = T$ , which is a contradiction.

Suppose that  $T = J_1$ . Then  $G = T$  and  $T_\alpha \cong D_{12}$ , so  $T_{\alpha\beta} \cong \mathbb{Z}_2^2$  for  $\beta \in \Gamma(\alpha)$ . It follows from the information given in the Atlas [8] that  $N_T(T_{\alpha\beta}) = \mathbb{Z}_2 \times (T_{\alpha\beta} : \mathbb{Z}_3) \cong \mathbb{Z}_2 \times A_4$ . Since all elements of order 6 of  $T$  are conjugate, all subgroups of  $T$  isomorphic to  $D_{12}$  are conjugate. Thus, we assume that  $T_\alpha$  is contained in a maximal subgroup  $M \cong \mathbb{Z}_2 \times A_5$ . Then  $N_M(T_{\alpha\beta}) \cong \mathbb{Z}_2^3$  is the Sylow 2-subgroup of  $N_T(T_{\alpha\beta})$ . Thus, there is no 2-element  $x \in N_T(T_{\alpha\beta})$  with  $\langle x, T_\alpha \rangle = T$ , which is a contradiction.

Assume that  $T = A_6$ . Then 12 divides  $|T_\alpha|$ , so  $T_\alpha \cong A_4$  or  $S_4$  by checking the subgroups of  $A_6$ . If  $T_\alpha \cong A_4$ , then  $T$  is transitive on  $V$ . Hence  $\Gamma$  is  $T$ -arc-transitive, and so  $A_4 \cong T_\alpha \geq S_3$  by Theorem 3.2, a contradiction. Thus  $T_\alpha \cong S_4$  and  $T$  has exactly two orbits on  $V$ , say  $U$  and  $W$ . Considering the possible permutation representations of  $A_6$  of degree 15, we may assume that each of  $U$  and  $W$  consists of either the 2-subsets of  $\Lambda := \{1, 2, 3, 4, 5, 6\}$ , or the partitions with part size 2 of  $\Lambda$ . Noting that, for  $\alpha \in U$ , the neighborhood  $\Gamma(\alpha)$  is a  $T_\alpha$ -orbit on  $W$ . Since  $|\Gamma(\alpha)| = 3$ , computation shows that, relabeling if necessary,  $U$  consists 2-subsets, and  $W$  consists of partitions, such that  $\alpha \in U$  is adjacent to  $\beta \in W$  if and only if  $\alpha$  is a part of  $\beta$ . Thus  $\Gamma$  is isomorphic to Tutte's 8-cage, and then part (1) of this lemma follows.

Now assume that  $T = \text{PSL}(2, p)$ , for a prime  $p \geq 5$ . Then  $G = \text{PSL}(2, p)$  or  $\text{PGL}(2, p)$ . Inspecting subgroups of  $G$  listed in [13, Chapter II, 8.27] and [3],  $G$  does not have subgroups isomorphic to  $S_4 \times S_2$ . Thus,  $G_\alpha$  is isomorphic to one of  $S_3$ ,  $D_{12}$ , and  $S_4$ . It follows that either  $T_\alpha = G_\alpha$ , or  $T_\alpha \cong S_3$  and  $G_\alpha \cong D_{12}$ .

First, let  $T_\alpha \cong S_3$ . Since  $|G : G_\alpha|$  is square-free, so is  $|T : T_\alpha|$ . Thus, 8 does not divide  $|T| = p(p^2 - 1)/2$ , and so  $p \equiv \pm 3 \pmod{8}$ . Since  $|T : T_\alpha|$  is even,  $T$  is transitive on  $V$ . Hence  $\Gamma$  can be written as a coset graph as in Example 3.5 (1).

Suppose now that  $T_\alpha = G_\alpha \cong D_{12}$ . Since  $|G : G_\alpha|$  is even and square-free, 8 divides  $|G|$  but 16 does not. Thus, either  $G = T = \text{PSL}(2, p)$ ,  $p \equiv \pm 7 \pmod{16}$  and  $\Gamma$  is isomorphic to a coset graph in Example 3.5 (2), or  $G = \text{PGL}(2, p)$ ,  $p \equiv \pm 3 \pmod{8}$  and  $\Gamma$  is isomorphic to a coset graph given in Example 3.6 (1).

In the case where  $T_\alpha = G_\alpha = S_4$ , the order  $|G|$  is divisible by 16 but not 32 since  $|G : G_\alpha|$  is even and square-free. Hence either  $G = T = \text{PSL}(2, p)$  with  $p \equiv \pm 15 \pmod{32}$  and  $\Gamma$  is isomorphic to the coset graph in Example 3.5 (3), or  $G = \text{PGL}(2, p)$  with  $p \equiv \pm 7 \pmod{16}$  and  $\Gamma$  is isomorphic to the coset graph in Example 3.6 (2). ■

Now we consider the case where  $G$  is not transitive on the arcs of  $\Gamma$ . Then  $\Gamma \cong \text{Cos}(G, G_\alpha\{x, y\}G_\alpha)$ , where  $x$  and  $y$  are 2-elements such that  $\langle x, y, G_\alpha \rangle = G$ ,  $\alpha^x, \alpha^y \in \Gamma(\alpha)$ ,  $x \in N_G(G_\alpha)$  with  $x^2 \in G_\alpha$ ,  $y \in N_G(G_{\alpha\alpha^y})$  with  $y^2 \in G_{\alpha\alpha^y}$ .

**Lemma 6.4.** *Assume that  $G$  is almost simple with socle  $\text{soc}(G) = T$  and  $\Gamma$  is not  $G$ -arc-transitive. Then  $T = \text{PSL}(2, p)$ , and either  $G_\alpha \cong \mathbb{Z}_2^2$ , or  $G_\alpha = T_\alpha \cong \mathbb{Z}_2$  or  $D_8$ ; moreover,  $\Gamma$  is isomorphic to a graph given in Examples 3.7 and 3.8.*

**Proof.** Since  $\Gamma$  is not  $G$ -arc-transitive and  $G$  is not regular,  $G_\alpha$  is a nontrivial 2-group. Then  $r^2$  is not a divisor of  $|G|$ , where  $r$  is an arbitrary odd prime. Checking the orders of finite simple groups,  $T = \text{soc}(G)$  is one of  $J_1$ ,  $\text{PSL}(2, p)$  for prime  $p \geq 5$ ,  $\text{PSL}(2, 2^f)$  with  $f \geq 4$ , and  $\text{Sz}(2^f)$  for odd  $f \geq 3$ .

Suppose that  $T = \text{PSL}(2, 2^f)$  with  $f \geq 4$  or  $\text{Sz}(2^f)$  for  $f \geq 3$ . Then any two distinct Sylow 2-subgroups of  $T$  intersect trivially, see [13, Chapter II, 8.5] and [21]. Now  $|T_\alpha| \geq 2^4$  and for  $\beta \in \Gamma(\alpha)$ , we have  $|T_\alpha : T_{\alpha\beta}| \leq 2$ , and hence  $T_{\alpha\beta} \neq 1$ . Thus,  $T_\alpha$  and  $T_\beta$  are contained in the same Sylow 2-subgroup  $Q$  of  $T$ . Since  $\Gamma$  is connected, it follows that  $T_\gamma \leq Q$  for all vertices  $\gamma$  of  $\Gamma$ . Hence,  $Q$  contains a nontrivial normal subgroup  $\langle T_\beta \mid \beta \in V\Gamma \rangle = \langle T_\alpha^g \mid g \in G \rangle$  of  $T$ , which is a contradiction.

Suppose that  $T = J_1$ . Then  $T = G$ , and since  $|T : T_\alpha|$  is even and square-free, we have  $T_\alpha \cong \mathbb{Z}_2^2$ . Let  $\beta \in \Gamma(\alpha)$  with  $T_{\alpha\beta} = \mathbb{Z}_2$ . Since  $\Gamma$  is connected,  $\langle T_\alpha, x, y \rangle = T$ , where  $x \in \mathbf{N}_T(T_\alpha)$  with  $x^2 \in T_\alpha$ , and  $y \in \mathbf{N}_T(T_{\alpha\beta})$  with  $y^2 \in T_{\alpha\beta}$ . By the Atlas [8],  $\mathbf{N}_T(T_{\alpha\beta}) \cong \mathbb{Z}_2 \times A_5$  and  $\mathbf{N}_T(T_\alpha) \cong \mathbb{Z}_2 \times A_4$ . Then  $x$  is contained in the unique Sylow 2-subgroup  $\langle T_\alpha, x \rangle$  of  $\mathbf{N}_T(T_\alpha)$ . Since  $T_{\alpha\beta} < \langle T_\alpha, x \rangle \cong \mathbb{Z}_2^3$ , we have  $x \in \langle T_\alpha, x \rangle < \mathbf{N}_T(T_{\alpha\beta})$ . Thus  $\langle x, y, G_\alpha \rangle \leq \mathbf{N}_T(T_{\alpha\beta}) \neq T$ , which is a contradiction.

Thus,  $T = \text{PSL}(2, p)$  for a prime  $p \geq 5$ . Then  $G = \text{PSL}(2, p)$  or  $\text{PGL}(2, p)$ , and a Sylow 2-subgroup of  $G$  is a dihedral group.

If  $|G_\alpha| = 2$ , then  $G_\alpha \cong \mathbb{Z}_2$ ,  $G = T = \text{PSL}(2, p)$  with  $p \equiv \pm 3 \pmod{8}$ , and  $\Gamma$  is isomorphic to a coset graph in Example 3.7 (1).

Assume that  $|G_\alpha| = 4$ . Then, by Lemma 3.3,  $G_\alpha$  is not cyclic, so  $G_\alpha \cong \mathbb{Z}_2^2$ . Hence either  $G = T = \text{PSL}(2, p)$  with  $p \equiv \pm 7 \pmod{16}$ , or  $G = \text{PGL}(2, p)$  with  $p \equiv \pm 3 \pmod{8}$ . For the former case,  $\Gamma$  is isomorphic to a coset graph in Example 3.7 (2). The latter case implies that  $T_\alpha \cong \mathbb{Z}_2$  or  $\mathbb{Z}_2^2$  depending on  $T$  is or not transitive on  $V$ , and so  $\Gamma$  is isomorphic to a coset graph in Example 3.7 (1) or 3.8 (1), respectively.

Finally, assume that  $G_\alpha = \langle a \rangle : \langle b \rangle \cong D_{2^e}$  for  $e \geq 3$ . Let  $\beta \in \Gamma(\alpha)$  with  $G_\alpha \neq G_\beta$ . Then  $G_{\alpha\beta}$  has index 2 in  $G_\alpha$ . If  $G_{\alpha\beta}$  contains a cyclic subgroup  $Z$  with  $|Z| \geq 4$ , then  $Z$  is characteristic in both  $G_\alpha$  and  $G_{\alpha\beta}$ , which contradicts with Lemma 3.3. Thus  $G_{\alpha\beta} \cong \mathbb{Z}_2^2$  and  $G_\alpha \cong D_8$ . Suppose that  $G_\alpha \neq T_\alpha$ . Then  $|T_\alpha| = 4$ ,  $G = \text{PGL}(2, p)$ , and  $T$  is transitive on  $V$ . Since  $T$  is not regular,  $T_{\alpha\beta} \cong \mathbb{Z}_2$ , and so  $G_{\alpha\beta} \not\leq T$ . Thus  $\mathbf{N}_G(G_{\alpha\beta}) \cong D_8$  by [3], so  $\mathbf{N}_G(G_{\alpha\beta}) = G_\alpha$ . Then there are no  $x \in \mathbf{N}_G(G_\alpha)$  and  $y \in \mathbf{N}_G(G_{\alpha\beta})$  such that  $\langle G_\alpha, x, y \rangle = G$ , a contradiction.

Therefore,  $G_\alpha = T_\alpha \cong D_8$ . Then either  $G = T = \text{PSL}(2, p)$  with  $p \equiv \pm 15 \pmod{32}$  and  $\Gamma$  is isomorphic to a coset graph in Example 3.7 (3), or  $G = \text{PGL}(2, p)$  with  $p \equiv \pm 7 \pmod{16}$  and  $\Gamma$  is isomorphic to a coset graph in Example 3.8 (2). ■

By Lemmas 6.3, 6.4, and their proofs, the next result determines some connected cubic Cayley graphs of square-free order which have insoluble automorphism groups.

**Corollary 6.5.** *Assume that  $T := \text{soc}(G) = \text{PSL}(2, p)$  for a prime  $p > 5$ . Then  $G$  contains no regular subgroups unless:*

- (1)  $G = \text{PGL}(2, 7)$ ,  $G$  has a regular subgroup  $R \cong D_{14}$ ,  $\mathbf{N}_G(R) = R : \mathbb{Z}_3$  and  $\Gamma$  is constructed as in Example 3.6 (2);
- (2)  $G = \text{PGL}(2, 7)$ ,  $G$  has a regular subgroup  $R \cong \mathbb{Z}_7 : \mathbb{Z}_6$ ,  $\mathbf{N}_G(R) = R$  and  $\Gamma$  is constructed as in Example 3.8 (2);
- (3)  $G = \text{PGL}(2, 11)$ ,  $G$  has a regular subgroup  $R \cong \mathbb{Z}_{11} : \mathbb{Z}_{10}$ ,  $\mathbf{N}_G(R) = R$  and  $\Gamma$  is constructed as in Example 3.6 (1);
- (4)  $G = \text{PGL}(2, 23)$ ,  $G$  has a regular subgroup  $R \cong \mathbb{Z}_{23} : \mathbb{Z}_{22}$ ,  $\mathbf{N}_G(R) = R$  and  $\Gamma$  is constructed as in Example 3.6 (2).

**Proof.** By Lemmas 6.3 and 6.4,  $T_\alpha$  (or  $G_\alpha$ ) and  $\Gamma$  are known and listed as follows:

$T_\alpha$	$G_\alpha$	$\Gamma$	$p$
$S_3$		3.5 (1)	5,11
$D_{12}$	$D_{12}$	3.5 (2), 3.6 (1)	5,7,11,23
$S_4$	$S_4$	3.5 (3), 3.6 (2)	7,23,47
$\mathbb{Z}_2$	$\mathbb{Z}_2$	3.7 (1)	None
	$\mathbb{Z}_2^2$	3.7 (1)–(2), 3.8 (1)	7
$D_8$	$D_8$	3.7 (3), 3.8 (2)	7

Suppose that  $G$  has a regular subgroup  $R$ . Then  $\Gamma$  is a Cayley graph and, since  $|G : T| \leq 2$ , we know that  $T$  contains a subgroup of order  $\frac{|R|}{2}$ . Thus  $T$  has a subgroup of square-free order  $\frac{|T|}{|T_\alpha|}$  or  $\frac{|T|}{2|T_\alpha|}$ , and such a subgroup has order divided by  $p$  as  $T_\alpha$  is a  $\{2, 3\}$ -group. Checking the subgroups of  $T$  (see [13], 8.27)), we conclude that  $p + 1$  divides  $|T_\alpha|$  or  $2|T_\alpha|$ . It follows that all possible  $p$  are listed at the last column of the above table. If  $p = 5$  then  $\Gamma$  is a 2-arc-transitive graph, and so  $\Gamma$  is the Petersen graph, which is not a Cayley graph. If  $p = 47$  then  $T_\alpha = G_\alpha \cong S_4$  and  $\Gamma$  is constructed as in Example 3.5 (3); however,  $G = T$  has no subgroup of order  $47 \cdot 46$ .

Assume that  $p = 7$ . Then  $G_\alpha \cong D_{12}$ ,  $S_4$ ,  $\mathbb{Z}_2^2$ , or  $D_8$ , and  $\Gamma$  is, respectively, constructed as in Example 3.5 (2), Example 3.6 (2), Example 3.7 (2), or Example 3.8 (2). Note that  $G$  has neither subgroups isomorphic to  $D_{12}$  and of square-free index, nor subgroups of order  $\frac{|G|}{4}$ . Then one of items (1) and (2) occurs.

Assume that  $p = 11$ . Then  $\Gamma$  is a 2-arc-transitive cubic graph of order 110. By [6], such a graph is isomorphic to a bipartite graph. It follows that  $T$  is not transitive on the vertices of  $\Gamma$ . Thus item (3) follows.

Finally, let  $p = 23$ . Then  $\Gamma$  is constructed as in Example 3.5 (2) or Example 3.6 (2). In this case, by the Atlas [8],  $G$  has no subgroups of order  $\frac{|G|}{12}$ , and then (4) follows. ■

Now we can determine the structure of  $G$  in the general case.

Let  $M$  be the soluble radical of  $G$  and let  $G^{(\infty)}$  be the smallest normal subgroup of  $G$  such that  $G/G^{(\infty)}$  is soluble. By Lemma 6.1,  $M$  has odd order and  $\Gamma$  is a cover of the quotient  $\Gamma_M$ , so  $\Gamma_M$  is cubic. Moreover,  $G/M$ , viewed as a transitive subgroup of  $\text{Aut}\Gamma_M$ , has trivial soluble radical. Then, by Lemmas 6.2, 6.3, and 6.4,  $G/M$  is almost simple with socle  $A_6$  or  $\text{PSL}(2, p)$ . Set  $\text{soc}(G/M) = Y/M$ . Then  $G/Y \cong (G/M)/(Y/M)$  is soluble, so  $G^{(\infty)} \leq Y$ . Thus  $Y = MG^{(\infty)}$ , and so  $G^{(\infty)}/(M \cap G^{(\infty)}) \cong MG^{(\infty)}/M = Y/M \cong A_6$  or  $\text{PSL}(2, p)$ .

On the other hand,  $\text{Aut}(M)$  is soluble as  $M$  has square-free order. Since  $G/\mathbf{C}_G(M) = \mathbf{N}_G(M)/\mathbf{C}_G(M)$  is isomorphic to a subgroup of  $\text{Aut}(M)$ , we have  $G^{(\infty)} \leq \mathbf{C}_G(M)$ . Then  $M \cap G^{(\infty)}$  is the center of  $G^{(\infty)}$ . Since  $M$  has odd order and  $3^3$  is not a divisor of  $|G|$ , we conclude that  $M \cap G^{(\infty)} = 1$  by checking the Schur multipliers of  $A_6$  and  $\text{PSL}(2, p)$ . Then  $Y = M \rtimes T$ , and so  $G = (M \rtimes T).O$ , where  $T = G^{(\infty)} = A_6$  or  $\text{PSL}(2, p)$ , and  $O$  lies in the outer automorphism group  $\text{Out}(T)$  of  $T$ .

**Lemma 6.6.** *Assume that  $G$  is insoluble. Then one of the following holds:*

- (1)  $G$  is almost simple with socle isomorphic to  $A_6$  or  $\text{PSL}(2, p)$ ;
- (2)  $\Gamma$  is not  $G$ -arc-transitive, and  $G = T:D_{2m}$  such that  $T = \text{PSL}(2, p)$ ,  $G_\alpha = T_\alpha \cong \mathbb{Z}_2^2$  is a Sylow 2-subgroup of  $T$ , and  $(|T|, m) = 1$ ;  $G$  contains no regular subgroups, and  $\Gamma$  can be constructed as in Construction 4.2.

**Proof.** Recall that  $G = (M \times T).O$ , where  $T = A_6$  or  $\text{PSL}(2, p)$ , and  $O \leq \text{Out}(T)$ .

If  $M = 1$ , then (1) follows from Lemmas 6.2, 6.3, and 6.4. Thus, we assume next that  $M \neq 1$ . Then  $m = |M| \geq 3$  is odd square-free.

Suppose that  $T$  has at most two orbits on  $V$ . Then  $M$  fixes one  $T$ -orbit  $U$ . By Lemma 6.1,  $M$  is semiregular and of odd square-free order. Then  $|M| \mid |U|$ , so  $|M| \mid |T|$ , and hence  $|M|^2 \mid |G|$ . Since  $|V| = |G : G_\alpha|$  is square-free for  $\alpha \in U$ , we have  $|M| \mid |G_\alpha|$ . Note that  $G_\alpha$  is either a 2-group or isomorphic to one of  $S_3$ ,  $D_{12}$ , and  $S_4$ . It follows that  $|M| = 3$  and  $3 \mid |G_\alpha|$ . Thus  $G_\alpha$  is 2-transitive on  $\Gamma(\alpha)$ , and so  $T_\alpha$  is transitive on  $\Gamma(\alpha)$  as  $T_\alpha$  is normal in  $G_\alpha$  and  $T$  is not semiregular on  $V$ ; in particular,  $3 \mid |T_\alpha|$ . Since  $|M| \mid |V|$  and  $|V| = |U|$  or  $2|U|$ , we know that 3 divides  $|U| = |T : T_\alpha|$ . Then  $3^2 \mid |T|$ , so  $3^3 \mid |G|$ , hence  $3^2 \mid |G_\alpha|$ , a contradiction. Thus  $T$  has at least three orbits on  $V$ .

Let  $K$  be the kernel of  $G$  acting on the  $T$ -orbits. Then, by Lemma 4.1,  $\Gamma_T \cong C_l$ ,  $G_\alpha = K_\alpha$  is a 2-group,  $l$  is even, and  $G/K = D_l$  acting regularly on  $T$ -orbits. Then  $M \cong KM/K \cong \mathbb{Z}_{\frac{l}{2}}$  and  $l = 2m$ . In particular,  $G$  is not transitive on the arcs of  $\Gamma$ , and so  $G/M$  is not transitive on the arcs of  $\Gamma_M$ . It follows from Lemma 6.4 that  $\text{soc}(G/M) \cong \text{PSL}(2, p)$ . Since  $K \geq T$  and  $|G/M| = \frac{|G|}{|M|} = \frac{l|K|}{m} = 2|K|$ , we have  $G/M \cong \text{PGL}(2, p)$  and  $K = T = \text{PSL}(2, p)$ . Clearly,  $\text{soc}(G/M)$  has two orbits on the vertices of  $\Gamma_M$ . By Lemma 6.4,  $(G/M)_\Delta \cong \mathbb{Z}_2^2$  or  $D_8$  for an  $M$ -orbit  $\Delta$ . Let  $\alpha \in \Delta$ . Then  $G_\Delta = MG_\alpha = MT_\alpha$ , and so  $T_\alpha \cong G_\Delta/M \cong (G/M)_\Delta \cong \mathbb{Z}_2^2$  or  $D_8$ . Since  $|V| = 2m|T : T_\alpha|$  is square-free,  $G_\alpha = T_\alpha$  is a Sylow 2-subgroup of  $T$  and  $m$  is coprime to  $|T|$ . Thus, we may assume that  $G = M:X$  with  $T < X \cong \text{PGL}(2, p)$ . Then  $N_G(G_\alpha) = MN_X(T_\alpha)$  and  $N_G(G_{\alpha\beta}) = MN_X(T_{\alpha\beta})$ , where  $\beta \in \Gamma(\alpha)$  with  $G_\alpha \neq G_\beta$ .

Suppose that  $G_\alpha = T_\alpha \cong D_8$ . Then  $N_X(G_\alpha) \cong D_{16}$ ,  $T_{\alpha\beta} = G_{\alpha\beta} \cong \mathbb{Z}_2^2$ ,  $S_4 \cong N_X(T_{\alpha\beta}) = N_T(T_{\alpha\beta})$ . Thus  $N_G(G_\alpha) = M:D_{16}$  and  $N_G(G_{\alpha\beta}) = M \times S_4$ . Then, for  $x \in N_G(G_\alpha)$  and  $y \in N_G(G_{\alpha\beta})$ , either  $\langle G_\alpha, x, y \rangle \leq M \times T$  or  $\langle G_\alpha, x, y \rangle \lesssim \text{PGL}(2, p)$ , which contradicts with the connectedness of  $\Gamma$ .

Assume that  $G_\alpha = T_\alpha \cong \mathbb{Z}_2^2$ . Then  $N_X(G_\alpha) \cong S_4$  and  $N_X(G_{\alpha\beta}) \cong D_{2(p-\varepsilon)}$ , where  $\varepsilon = \pm 1$  such that  $4 \parallel p - \varepsilon$ . Note that  $G_\alpha \leq N_X(G_{\alpha\beta})$ . Take an involution  $b \in N_X(G_{\alpha\beta})$  with  $G_\alpha : \langle b \rangle \cong D_8$ . Then  $b \in X \setminus T$ ,  $M : \langle b \rangle \cong D_{2m}$ ,  $N_G(G_\alpha) = (M \times N_T(T_\alpha)) \langle b \rangle$  and  $N_G(G_{\alpha\beta}) = M \times N_T(T_{\alpha\beta}) \langle b \rangle$ . Thus  $\Gamma$  can be constructed as in Construction 4.2.

Suppose that  $G$  has a regular subgroup. Then, since  $|G : MT| = 2$ , we know that  $MT = M \times T$  contains a subgroup of order  $\frac{|G:G_\alpha|}{2} = \frac{|MT|}{4}$ . Thus  $T$  has a subgroup of index 4, which is impossible as  $T$  is simple. Then the result follows. ■

## 7. PROOF OF THEOREM 1.1

Let  $\Gamma$  be a connected vertex-transitive cubic graph of square-free order  $2n$ .

If  $\text{Aut}\Gamma$  is insoluble then  $\Gamma$  is known as in parts (2)–(4) of Theorem 1.1 by the argument in Section 6. To complete the proof, we first determine the Cayley graphs which have insoluble automorphism groups. Assume that  $\text{Aut}\Gamma$  is insoluble and has a regular subgroup  $G$ . By Corollary 6.5 and Lemma 6.6 (2), either

- (i)  $\text{Aut}\Gamma = \text{PGL}(2, 7)$ ,  $G = \langle a \rangle : \langle b \rangle \cong D_{14}$ , and  $N_{\text{Aut}\Gamma}(R) = R : \mathbb{Z}_3$ ; or
- (ii)  $\text{Aut}\Gamma = \text{PGL}(2, p)$ ,  $G = \langle a \rangle : \langle b \rangle \cong \mathbb{Z}_p : \mathbb{Z}_{p-1}$ , and  $N_{\text{Aut}\Gamma}(R) = R$ , where  $p \in \{7, 11, 23\}$ .

For (i), by Lemma 2.3 (3),  $\Gamma \cong \text{Cay}(G, \{ab, a^3b, b\})$  or  $\text{Cay}(R, \{ab, a^5b, b\})$ . Verified by **Magma**,  $\text{Cay}(R, \{ab, a^3b, b\}) \cong \text{Cay}(R, \{ab, a^5b, b\})$ , so Line 1 of Table I occurs. For (ii), by Lemma 2.5,  $\Gamma \cong \text{Cay}(G, \{ab^k, (ab^k)^{-1}, b^l\})$  with  $a^{b^{\frac{p-1}{2}}} = a^{-1}$ ,  $0 < k < \frac{p-1}{2}$  and  $(k, \frac{p-1}{2}) = 1$ . Then, verified by **Magma**, one of Lines 2, 4, and 5 of Table I occurs.

Now assume that  $\text{Aut}\Gamma$  is soluble. Then either  $\Gamma$  is a Cayley graph or a generalized Petersen graph by the argument in Section 5, and hence  $\Gamma$  is known by the argument in Section 2. Assume that  $\Gamma \cong \mathbf{P}(n, r)$  is a generalized Petersen graph, where  $1 \leq r < \frac{n}{2}$ . If  $r^2 \equiv 1 \pmod{n}$  then, by [11],  $\text{Aut}\mathbf{P}(n, r) \cong \mathbb{Z}_n : \mathbb{Z}_2^2$  contains a regular subgroup described as in (i), and it is easily shown that  $\mathbf{P}(n, r)$  is neither a circulant nor a dihedrant unless  $r = 1$ . For  $r^2 \equiv -1 \pmod{n}$ , again by [11], either  $\text{Aut}\mathbf{P}(n, r) \cong \mathbb{Z}_n : \mathbb{Z}_4$  or  $(n, r) = (5, 2)$  and  $\Gamma$  is the Petersen graph; moreover, in this case,  $\Gamma$  is not isomorphic to a Cayley graph. Then one of Theorem 1.1 (i) and (vii) occurs.

Therefore, we assume next that  $\Gamma = \text{Cay}(G, S)$  is a Cayley graph. If  $G$  has a subgroup isomorphic to  $\mathbb{Z}_n$  then  $G \cong \mathbb{Z}_n : \mathbb{Z}_2$ , hence  $\text{Aut}\Gamma = \bar{G} : \text{Aut}(G, S)$  and one of (i)–(v) occurs by Lemmas 2.2–2.5, Corollary 2.4 and the argument in Section 5.

Suppose that  $G$  has no subgroups isomorphic to  $\mathbb{Z}_n$ . By Lemmas 2.1 and 2.5, we may assume that  $n > 3$ ,  $\Gamma = \text{Cay}(G, S_k)$  and  $\text{Aut}(G, S_k) = 1$ , where  $G = \langle c \rangle \times (\langle a \rangle : \langle b \rangle)$ ,  $o(b) = 2l > 2$ ,  $\mathbf{Z}(G) = \langle c \rangle$ ,  $G' = \langle a \rangle$ ,  $a^{b^l} = a^{-1}$ ,  $S_k = \{cab^k, (cab^k)^{-1}, b^l\}$ ,  $1 < k < l$  and  $(k, l) = 1$ . Then, by the argument in Section 5, either  $\text{Aut}\Gamma = \bar{G}$  or  $\text{Aut}\Gamma \cong \mathbb{Z}_n : \mathbb{Z}_6 \cong D_{2n} : \mathbb{Z}_3$ . We next show Theorem 1.1 (vi) occurs, it suffices to show that  $\text{Aut}\Gamma \cong \mathbb{Z}_n : \mathbb{Z}_6$  if and only if  $G$  and  $k$  are described as in Line 3 of Table I.

Suppose that  $G = \langle a \rangle : \langle b \rangle$  with  $o(b) = 6$  and  $a^b = a^t$  such that  $t^2 - t + 1 \equiv 0 \pmod{n}$ . Let  $\Gamma = \text{Cay}(G, S)$ , where  $S = \{ab, (ab)^{-1}, b^3\}$ . Define a map

$$\pi : G \rightarrow, a^i b^j \mapsto \begin{cases} a^{it^2}, & \text{if } j \equiv 0 \pmod{6}; \\ a^{it^2-t+1} b^2, & \text{if } j \equiv 2 \pmod{6}; \\ a^{it^2-t} b^4, & \text{if } j \equiv 4 \pmod{6}; \\ a^{-it} b^5, & \text{if } j \equiv 1 \pmod{6}; \\ a^{-it+1} b, & \text{if } j \equiv 3 \pmod{6}; \\ a^{-it-t+1} b^3, & \text{if } j \equiv 5 \pmod{6}. \end{cases}$$

It is easily shown  $\pi$  is an automorphism of  $\Gamma$  and fixes the vertex 1. Note that all Cayley graphs with insoluble automorphism groups are known, whose order is either 42 or not divisible by 3. If  $|G| = 42$  then, verified by **Magma**,  $\text{Aut}\Gamma$  is soluble and has order 126. Thus, we conclude that  $\text{Aut}\Gamma$  is soluble. By the argument in Section 5, we conclude that  $\text{Aut}\Gamma \cong \mathbb{Z}_n : \mathbb{Z}_6$ .

Suppose now that  $\text{Aut}\Gamma \cong \mathbb{Z}_n : \mathbb{Z}_6$ . Then  $\text{Aut}\Gamma$  has a unique  $\{2, 3\}'$ -Hall subgroup  $L$ . Clearly,  $L$  is cyclic and normal in  $\text{Aut}\Gamma$ . Consider the subgroup  $X := L\bar{G}$  of  $\text{Aut}\Gamma$ . Since  $X$  is transitive on the vertices of  $\Gamma$ , we have  $X = \bar{G}X_\alpha$  for some vertex  $\alpha$ . Then  $\frac{|L||G|}{|L \cap G|} = |L\bar{G}| = |X| = |G||X_\alpha| = |G|$  or  $3|G|$ , yielding  $L < \bar{G}$ . Thus  $L$  is a cyclic normal subgroup of  $\bar{G}$ . Let  $N$  be the Fitting subgroup of  $\bar{G}$ . Then  $L \leq N$ . Since  $\bar{G}$  has square-free order,  $N$  is cyclic. It is easily shown that  $N = \langle \bar{c} \rangle \times \langle \bar{a} \rangle$ . Then  $2l = |\bar{G} : N|$  divides  $|\bar{G} : L|$ , so  $|\bar{G} : L| \geq 2l \geq 6$ . Note that  $L$  is a  $\{2, 3\}'$ -Hall subgroup of  $\bar{G}$ . Thus  $|\bar{G} : L|$  divides 6, and so  $2l$  divides 6. Thus  $2l = 6$  as  $l > 1$ , and hence  $L = N$ . Since  $0 < k < l = 3$ , we have  $k = 1$  or 2.

Consider the normal quotient graph  $\Gamma_N$ . We know that  $\Gamma_N \cong \text{Cay}(\langle b \rangle, \{b^k, b^{-k}, b^3\})$ . Then either  $\Gamma_N \cong K_{3,3}$  for  $k = 1$ , or  $\Gamma_N \cong P(3, 1)$  for  $k = 2$ . Since  $N$  is normal in  $\text{Aut}\Gamma$  and  $\Gamma$  is arc-transitive,  $\Gamma_N$  is also arc-transitive. It follows that  $k = 1$ .

By Lemma 5.3,  $\text{Aut}\Gamma$  has a normal regular subgroup  $R \cong D_{2n}$ . Note that each Sylow 2-subgroup of  $\text{Aut}\Gamma \cong \mathbb{Z}_n : \mathbb{Z}_6$  has order 2. It follows that all involutions in  $\text{Aut}\Gamma$  are conjugate. Thus we may choose  $R$  such that  $\bar{b}^3 \in R$ . Recalling  $L = N = \langle \bar{c}, \bar{a} \rangle$  is the  $\{2, 3\}'$ -Hall subgroup of  $\text{Aut}\Gamma$ , we have  $N = \langle \bar{c}, \bar{a} \rangle < R$ . Then  $\bar{c}^{\bar{b}^3} = \bar{c}^{-1}$ , yielding  $o(c) = o(\bar{c}) = 1$  as  $\bar{c}\bar{b} = \bar{b}\bar{c}$ . Thus  $o(a) = \frac{n}{3}$  and  $\bar{G} \cong G = \langle a, b \rangle$  has trivial center. Moreover,  $R = \langle \bar{a}z, \bar{b}^3 \rangle$  for some  $z$  with  $o(z) = 3$  and  $z\bar{a} = \bar{a}z$ . It is easily shown that  $\langle \bar{a}z \rangle \cap \langle \bar{b} \rangle \leq \mathbf{Z}(\bar{G})$ . Then  $\langle \bar{a}z \rangle \cap \langle \bar{b} \rangle = 1$ , and so  $\text{Aut}\Gamma = \langle \bar{a}z \rangle : \langle \bar{b} \rangle = R : \langle \bar{b}^2 \rangle$ .

Assume that  $\theta \in \text{Aut}\Gamma$  has order 3. Note that  $\text{Aut}\Gamma$  has an abelian Sylow 3-subgroup  $\langle z, \bar{b}^2 \rangle$ . Then  $\theta \in \langle z, \bar{b}^2 \rangle^{\bar{a}^i}$  for some  $i$ . Assume further that  $\theta$  fixes the vertex 1 of  $\Gamma$ . Then, replacing  $z$  by  $z^{-1}$  if necessary, we may set  $\theta = z\bar{g}$  for  $g = a^{-i}b^{\pm 2}a^i$ . Thus  $1 = 1^\theta = 1^z g$ , and so  $1^z = g^{-1}$ . Since  $z\bar{g} = \bar{g}z$ , we have  $1 = 1^\theta = 1^{\bar{g}z} = g^z$ , and so  $1^{z^{-1}} = g$ . Let  $a^b = a^r$  for some  $r$  coprime to  $\frac{n}{3}$ . Then  $r^6 \equiv 1 \pmod{\frac{n}{3}}$  and  $r^3 \equiv -1 \pmod{\frac{n}{3}}$ . Thus  $(b^3)^\theta = 1^{\bar{b}^3} z\bar{g} = 1^{z^{-1}\bar{b}^3} \bar{g} = g\bar{b}^3 g = a^{-i(r+1)^2} b$  or  $a^{-i(r^2-1)^2} b^{-1}$ . Since  $\Gamma$  is arc-transitive,  $\langle \theta \rangle$  is transitive on  $\{ab, (ab)^{-1}, b^3\}$ . Then  $(b^3)^\theta = ab$  or  $(ab)^{-1}$ . Therefore, either  $a^{-i(r+1)^2} b = ab$  or  $a^{-i(r^2-1)^2} b^{-1} = (ab)^{-1} = a^{-r} b^{-1}$ . Then  $-i(r+1)^2 \equiv 1 \pmod{\frac{n}{3}}$  or  $-i(r^2-1)^2 \equiv -r \pmod{\frac{n}{3}}$ , it follows that  $(r+1, \frac{n}{3}) = 1$ . Since  $r^3 \equiv -1 \pmod{\frac{n}{3}}$ , we have  $r^2 - r + 1 \equiv 0 \pmod{\frac{n}{3}}$ .

Since  $\langle \bar{a}z \rangle$  is normal in  $\text{Aut}\Gamma$ , we set  $(\bar{a}z)^{\bar{b}} = (\bar{a}z)^t$  for some  $t$  coprime to  $n$ . Then  $(\bar{a}z)^{t^3} = (\bar{a}z)^{\bar{b}^3} = \bar{a}^{\bar{b}^3} z^{\bar{b}^3} = \bar{a}^{-1} z^{-1} = (\bar{a}z)^{-1}$ , so  $t^3 \equiv -1 \pmod{n}$ , hence  $t^3 \equiv -1 \pmod{\frac{n}{3}}$ . Note that  $\bar{a}' z' = (\bar{a}z)^t = (\bar{a}z)^{\bar{b}} = \bar{a}^{\bar{b}} z^{\bar{b}} = \bar{a}' z^{\bar{b}^4 \bar{b}^3} = \bar{a}' z^{-1}$ . It follows that  $t \equiv r \pmod{\frac{n}{3}}$  and  $t \equiv -1 \pmod{3}$ . Since  $t \equiv -1 \pmod{3}$ , we know that  $3 \mid (t^2 - t + 1)$ . Since  $r^2 - r + 1 \equiv 0 \pmod{\frac{n}{3}}$  and  $t \equiv r \pmod{\frac{n}{3}}$ , we have  $t^2 - t + 1 \equiv 0 \pmod{\frac{n}{3}}$ . Then, since  $(3, \frac{n}{3}) = 1$ , we have  $t^2 - t + 1 \equiv 0 \pmod{n}$ . Thus Theorem 1.1 (vi) occurs. This completes the proof.

## REFERENCES

- [1] B. Alspach and R. J. Sutcliffe, Vertex-transitive graphs of order  $2p$ , *Ann New York Acad Sci* 319 (1979), 18–27.
- [2] N. L. Biggs, *Algebraic Graph Theory*, 2nd edn., Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1986.
- [3] P. J. Cameron, G. R. Omidi, and B. Tayfeh-Rezaie, 3-Design from  $\text{PGL}(2, q)$ , *The Electronic J Combin* 13 (2006), #R50.
- [4] C. Y. Chao, On the classification of symmetric graphs with a prime number of vertices, *Trans Amer Math Soc* 158 (1971), 247–256.
- [5] Y. Cheng and J. Oxley, On weakly symmetric graphs of order twice a prime, *J Combin Theory Ser B* 42 (1987), 196–211.
- [6] M. Conder and P. Dobcsányi, Trivalent symmetric graphs on up to 768 vertices, *J Combin Math Combin Comput* 40 (2002), 41–63.
- [7] M. D. Conder, C. H Li, and C. E. Praeger, On the Weiss conjecture for finite locally primitive graphs, *Pro Edinburgh Math Soc* 43 (2000), 129–138.

- [8] J. H. Conway, R. T. Curtis, S. P. Noton, R. A. Parker, and R. A. Wilson, *Atlas of Finite Groups*, Clarendon Press, Oxford, 1985.
- [9] Y. Q. Feng, J. H. Kwak, X. Y. Wang, and J. X. Zhou, Tetravalent half-arc-transitive graphs of order  $2pq$ , *J Algebraic Comb* 33 (2011), 543–553.
- [10] Y. Q. Feng and Y. T. Li, One-regular graphs of square-free order of prime valency, *European J Combin* 32 (2011), 265–275.
- [11] R. Frucht, J. E. Graver, and M. E. Watkins, The groups of the generalized Petersen graphs, *Proc Cambridge Philos Soc* 70 (1971), 211–218.
- [12] C. D. Godsil, On the full automorphism group of a graph, *Combinatorica* 1 (1981), 243–256.
- [13] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, Berlin, 1967.
- [14] C. H. Li, Z. Liu, and Z. P. Lu, Tetravalent edge-transitive Cayley graphs of square free order, *Discrete Math* 312 (2012), 1952–1967.
- [15] C. H. Li, D. Marušič, and J. Morris, Classifying arc-transitive circulants of square-free order, *J Algebraic Combin* 14 (2001), 145–151.
- [16] C. H. Li, S. J. Song, and D. J. Wang, A characterization of metacirculants, *J Combin Theory A* 120 (2013), 39–48.
- [17] Y. T. Li and Y. Q. Feng, Pentavalent one-regular graphs of square-free order, *Algebra Colloq* 17 (2010), 515–524.
- [18] D. Marušič and R. Scapellato, Classifying vertex-transitive graphs whose order is a product of two primes, *Combinatorica* 14(2) (1994), 187–201.
- [19] C. E. Praeger, R. J. Wang, and M. Y. Xu, Symmetric graphs of order a product of two distinct primes, *J Combin Theory Ser B* 58 (1993), 299–318.
- [20] C. E. Praeger and M. Y. Xu, Vertex-primitive graphs of order a product of two distinct primes, *J Combin Theory Ser B* 59 (1993), 245–266.
- [21] M. Suzuki, On a class of doubly transitive groups, *Ann Math* 75(1) (1962), 105–145.
- [22] J. Turner, Point-symmetric graphs with a prime number of points, *J Combin Theory* 3 (1967), 136–145.
- [23] J. X. Zhou and Y. Q. Feng, Cubic vertex-transitive graphs of order  $2pq$ , *J Graph Theory* 65 (2010), 285–302.
- [24] J. X. Zhou and Y. Q. Feng, Cubic one-regular graphs of order twice a square-free integer, *Sci China Ser A: Math* 51 (2008), 1093–1100.