

Miscellanea

Multivariate sign-based high-dimensional tests for sphericity

BY CHANGLIANG ZOU, LIUHUA PENG, LONG FENG AND ZHAOJUN WANG

School of Mathematical Sciences, Nankai University, Tianjin, China

nk.chlzou@gmail.com allenberify@yahoo.com.cn fnankai@126.com zjwang@nankai.edu.cn

SUMMARY

This article concerns tests for sphericity in cases where the data dimension is larger than the sample size. The existing multivariate sign-based procedure (Hallin & Paindaveine, 2006) for sphericity is not robust with respect to high dimensionality, producing tests with Type I error rates that are much larger than the nominal levels. This is mainly due to bias from estimating the location parameter. We develop a correction that makes the existing test statistic robust with respect to high dimensionality, and show that the proposed test statistic is asymptotically normal under elliptical distributions. The proposed method allows the dimensionality to increase as the square of the sample size. Simulations demonstrate that it has good size and power in a wide range of settings.

Some key words: Asymptotic normality; Large p , small n ; Spatial median; Spatial sign; Sphericity test.

1. INTRODUCTION

High-dimensional data have dimension p that increases to infinity as the number of observations n tends to infinity. Traditional statistical methods may fail in this situation, since they are often based on assuming that p remains constant. Overcoming this challenge requires new research on the properties of traditional methods (see, e.g., Chen et al., 2009; Hjort et al., 2009) as well as the development of new statistical approaches to deal with high-dimensional data. Some methods were proposed by Chen & Qin (2010) for a two-sample test for means, by Ledoit & Wolf (2002), Schott (2005) and Chen et al. (2010) for testing a specific covariance structure, and by Tang & Leng (2010) and the references therein for variable selection.

Sphericity assumptions play a key role in a number of statistical problems. The need to test for sphericity arises from various applications, such as geostatistics, paleomagnetic studies, animal navigation, astronomy, wind direction data analysis and microarray analysis; see Tyler (1987), Baringhaus (1991), Marden & Gao (2002) and Sirkiä et al. (2009) for further information. Given the interest in both high-dimensional data and sphericity testing, the asymptotic and finite-sample properties of sphericity tests in the high-dimensional setting merit careful investigation.

Because of its importance in applications, sphericity testing has a long history and has generated a considerable body of literature, which we review only very briefly here. Let X_1, \dots, X_n denote a p -dimensional sample of size n . The distribution of a p -dimensional random vector X is said to be spherical if for some $\theta \in \mathbb{R}^p$, the distribution of $X - \theta$ is invariant under orthogonal transformations. For multinormal variables, sphericity is equivalent to Σ , the covariance matrix of X , being proportional to the identity matrix I_p . Thus approaches based on the covariance matrix, such as the likelihood ratio test (Mauchly, 1940), are popular. John (1971, 1972) considered the testing problem in the normal distribution case and showed that the statistic

$$Q_J = \frac{np^2}{2} \operatorname{tr} \left\{ \frac{S}{\operatorname{tr}(S)} - \frac{1}{p} I_p \right\}^2 \quad (1)$$

provides the locally most powerful invariant test for sphericity under the multivariate normality assumption; here S is the sample covariance matrix and I_p is the identity matrix. This test is valid only under multivariate

normality. In the wider elliptical model, one can use a modification of John's test statistic, $Q_M = Q_J/\hat{\kappa}$, where $\hat{\kappa}$ is the estimated kurtosis based on the standardized fourth moment of the marginal distribution (Muirhead & Waternaux, 1980).

Ledoit & Wolf (2002) evaluated Q_J in the case where the dimension p increases at the same rate as n , so that $p/n \rightarrow c$ for some finite c . Chen et al. (2010) developed a high-dimensional test based on Q_J , and showed that their proposed test statistic is asymptotically normal by assuming that the data have the structure $X_i = \Gamma Z_i + \mu$ where Γ is a $p \times m$ matrix, with $m \geq p$, and $Z_i = (Z_{i1}, \dots, Z_{im})^\top$ is a random vector such that

$$\begin{aligned} E(Z_i) &= 0, \quad \text{var}(Z_i) = I_m, \quad E(Z_{ii}^{4k}) = m_{4k} \in (0, \infty), \\ E(Z_{ik_1}^{\alpha_1} Z_{ik_2}^{\alpha_2} \cdots Z_{ik_q}^{\alpha_q}) &= E(Z_{ik_1}^{\alpha_1}) E(Z_{ik_2}^{\alpha_2}) \cdots E(Z_{ik_q}^{\alpha_q}) \end{aligned} \quad (2)$$

whenever $\sum_{k=1}^q \alpha_k \leq 4k$. Here k, k_1, \dots, k_q are positive integers. The data structure (2) generates a rich collection of X_i from Z_i with a given covariance Σ . For example, distributions in the so-called independent component model lie in the family given by (2); see Example 2.6 in Oja (2010) and the references therein. It is difficult, however, to justify this model. The condition that power transformations of different components of Z_i be uncorrelated is almost equivalent to saying that Z_{i1}, \dots, Z_{im} are independent, and is therefore not easily met in practice. For instance, it can be verified that a random vector from the multivariate t distribution or mixtures of multivariate normal distributions does not satisfy (2). Moreover, the statistical performance of this test would be degraded when the nonnormality is severe, especially for heavy-tailed distributions; see § 3. This motivates us to consider using multivariate sign- and/or rank-based covariance matrices to construct robust tests for sphericity.

Such an approach has been adopted by Tyler (1987), Ghosh & Sengupta (2001), Marden & Gao (2002), Hallin & Paindaveine (2006) and Sirkiä et al. (2009), among others. Most of the tests proposed by these researchers are based on the signs and ranks of the norms of the observations centred at θ , with test statistics that have structures similar to Q_J . These statistics are distribution-free, or asymptotically so, under sphericity and elliptical distributional assumptions. Hallin & Paindaveine (2006) and Oja (2010) provide nice overviews of this topic. Among these tests, one that is entirely based on multivariate signs, also called the spatial-sign test by some authors, is of particular interest due to its simplicity and effectiveness, and has been discussed in detail by Marden & Gao (2002), Hallin & Paindaveine (2006) and Sirkiä et al. (2009). In the present paper, we focus on this type of test. The existing calibration method is not robust with respect to high dimensionality in the sense that it would produce tests with Type I error rates much larger than the nominal levels. This is mainly due to bias arising from estimating the location parameter. In the next section, we develop a bias correction to the existing test statistic that makes it robust with respect to high dimensionality. We show that the proposed test statistic is asymptotically normal for elliptical distributions. Simulation comparisons show that our procedure has good size and power for a wide range of dimensions, sample sizes and distributions. Finite-sample studies also demonstrate that the proposed method works reasonably well when the underlying distribution is not elliptical, especially for observations from the data structure (2). All the proofs are given in the online Supplementary Material.

2. HIGH-DIMENSIONAL TEST FOR SPHERICITY USING MULTIVARIATE SIGNS

2.1. Inference based on the sign covariance matrix

Let X_1, \dots, X_n be a random sample from a p -variate elliptical distribution with density function $\det(\Sigma_p)^{-1/2} g_p\{\|\Sigma_p^{-1/2}(X - \theta_p)\|\}$, where $\|X\| = (X^\top X)^{1/2}$ is the Euclidean length of the vector X , θ_p is the symmetry centre and Σ_p is a positive-definite symmetric $p \times p$ scatter matrix. The matrix Σ_p that describes the covariances between the p variables can be expressed as $\Sigma_p = \sigma_p \Lambda_p$, where $\sigma_p = \sigma(\Sigma_p)$ is a scale parameter and $\Lambda_p = \sigma_p^{-1} \Sigma_p$ is a shape matrix. The scale parameter is assumed to satisfy $\sigma(I_p) = 1$ and $\sigma(a\Sigma_p) = a\sigma(\Sigma_p)$ for all $a > 0$. We wish to test the null hypothesis $H_0: \Sigma_p = \sigma I_p$. Under the assumption of ellipticity, finite second-order moments need not exist, and sphericity is equivalent to $\Lambda_p = I_p$. If one wishes to test the hypothesis $H_0: \Lambda_p = V$, one can use the standardized observations $V^{-1/2} X_i$

instead of the original ones. In the following, we assume that the shape matrix is standardized so that $\text{tr}(\Lambda_p) = p$.

The multivariate sign function is defined as $U(X) = \|X\|^{-1}XI$ ($X \neq 0$). The observed signs for the X_i are $U_i = U(X_i - \theta_p)$. Accordingly, the sign covariance matrix is defined by $\Omega_{n,p} = n^{-1} \sum_{i=1}^n U_i U_i^T$ (Hallin & Paindaveine, 2006). Under the null hypothesis, we have $E(\Omega_{n,p}) = I_p/p$. The sign test statistic can be defined by mimicking John's test statistic (1) with $\Omega_{n,p}$ in place of S (Hallin & Paindaveine, 2006; Sirkiä et al., 2009):

$$Q_S = p \text{tr}\{\text{tr}(\Omega_{n,p})^{-1} \Omega_{n,p} - p^{-1} I_p\}^2 = p \text{tr}(\Omega_{n,p} - p^{-1} I_p)^2.$$

It can be shown that when p is fixed, under the null hypothesis one has

$$n(p + 2)Q_S/2 \rightarrow \chi_{(p+2)(p-1)/2}^2 \tag{3}$$

in distribution as $n \rightarrow \infty$; see Hallin & Paindaveine (2006) for the proof.

In high-dimensional settings, p diverges to infinity as $n \rightarrow \infty$, so χ_p^2 is asymptotically normal with mean p and variance $2p$, and we might expect that

$$\text{var}(Q_S)^{-1/2} \{Q_S - E(Q_S)\} \rightarrow N(0, 1) \tag{4}$$

in distribution as $n \rightarrow \infty$ and $p \rightarrow \infty$. In what follows, we will show that the above convergence in law is essentially correct under mild conditions. However, the main impact of high dimensionality on the validity of the sign-based test does not stem from the difference between the two asymptotic calibrations of Q_S , (3) and (4). In the above discussion, the true location parameter θ_p is used in the definition of the sign vector, but in practice θ_p must usually be replaced by an estimator $\hat{\theta}_{n,p}$. Any root- n consistent estimator would be adequate, but in the literature the rotation-equivariant spatial median (Möttönen & Oja, 1995), which minimizes the criterion function $L(\theta) = \sum_{i=1}^n \|X_i - \theta\|$, is usually recommended. Taking the gradient of the objective function, one sees that $\hat{\theta}_{n,p}$ is the solution to the equation

$$\sum_{i=1}^n U(X_i - \theta) = 0.$$

When p is fixed, replacing θ_p with $\hat{\theta}_{n,p}$ does not affect the asymptotic properties of Q_S . However, as we shall show in the next section, this substitution would yield a bias term which is not negligible when $n/p = O(1)$. Even worse, when $n/p = o(1)$, the test based on (3) or (4) would have asymptotic size 1 under H_0 . We will propose a simple remedy for this problem.

2.2. A bias-corrected sign-based procedure

The test statistic Q_S can be rewritten as

$$Q_S = \frac{p}{n} + \frac{n(n-1)}{n^2} \frac{p}{n(n-1)} \sum_{i \neq j} (U_i^T U_j)^2 - 1,$$

so we consider the modified test statistic

$$Q'_S = \frac{p}{n(n-1)} \sum_{i \neq j} (U_i^T U_j)^2 - 1.$$

Upon substituting the spatial median $\hat{\theta}_{n,p}$ into U_i , the test statistic becomes

$$\tilde{Q} = \frac{p}{n(n-1)} \sum_{i \neq j} (\hat{U}_i^T \hat{U}_j)^2 - 1,$$

where $\hat{U}_i = U(X_i - \hat{\theta}_{n,p})$.

PROPOSITION 1. *The test statistic \tilde{Q} is invariant under rotations.*

The value of \tilde{Q} remains unchanged for $Z_i = aOX_i + c$ where a is a constant, c is a vector of constants and O is an orthogonal matrix. Thus, without loss of generality, we assume that $\theta_p = 0$ in what follows. It is easy to see that $E(\tilde{Q}_S) = 0$ under H_0 . However, in high-dimensional settings, $E(\tilde{Q})$ is not negligible with respect to $\text{var}^{1/2}(\tilde{Q})$. Before proceeding, we state a necessary assumption which is required throughout this paper. Let $R_i = \|X_i - \theta_p\|$.

Assumption 1. The moments $E(R_i^{-k})$ for $k = 1, \dots, 4$ exist for large enough p ; and, for $k = 2, 3, 4$, $E(R_i^{-k})/E(R_i^{-1})^k \rightarrow d_k \in [1, \infty)$ as $p \rightarrow \infty$, where the d_k are constants.

This assumption ensures the validity of the second-order expansions that we use and the existence of our bias-correction term. The moments $E(R_i^{-k})$ may not exist for a fixed p ; for example, for standard multivariate normal and t distributions, $E(R_i^{-2})$ is equal to $1/(p - 2)$ and thus the second moment exists only when $p > 3$. In the Supplementary Material, we verify this assumption for three commonly used elliptical distributions: the multivariate normal distribution, the multivariate t distribution, and mixtures of multivariate normal distributions. We also formulate this assumption using the g_p that fixes the distribution of the modulus R_i . The existence of $E(R_i^{-k})$ is guaranteed if $r^{p-1-k}g_p(r)$ is bounded for $r \in (0, \epsilon)$.

We define

$$\begin{aligned} \delta_{n,p} = & \frac{1}{n^2} \left(2 - \frac{2E(R_i^{-2})}{E(R_i^{-1})^2} + \left[\frac{E(R_i^{-2})}{E(R_i^{-1})^2} \right]^2 \right) \\ & + \frac{1}{n^3} \left[\frac{8E(R_i^{-2})}{E(R_i^{-1})^2} - 6 \left\{ \frac{E(R_i^{-2})}{E(R_i^{-1})^2} \right\}^2 + \frac{2E(R_i^{-2})E(R_i^{-3})}{E(R_i^{-1})^5} - \frac{2E(R_i^{-3})}{E(R_i^{-1})^3} \right] \end{aligned} \tag{5}$$

and $\tilde{\sigma}_0^2 = 4(p - 1)/\{n(n - 1)(p + 2)\}$. From the proof of Theorem 1 in the Supplementary Material, we know that

$$E(\tilde{Q}) = p\delta_{n,p} + o(n^{-1}), \quad \text{var}(\tilde{Q}) = \tilde{\sigma}_0^2 + o(n^{-2}).$$

Clearly, if $n < p$, using the normal calibration for \tilde{Q} as in (4) would result in a bias term which cannot be ignored, producing Type I error rates much larger than nominal levels. Hence, the key idea underpinning our method is to correct this bias through approximating $E(\tilde{Q})$. The following theorem establishes the asymptotic null distribution of \tilde{Q} .

THEOREM 1. *Under H_0 and Assumption 1, if $p = O(n^2)$, then $(\tilde{Q} - p\delta_{n,p})/\tilde{\sigma}_0 \rightarrow N(0, 1)$ in distribution as $p \rightarrow \infty$ and $n \rightarrow \infty$.*

The unknown quantities in $\delta_{n,p}$ are $E(R_i^{-2})/\{E(R_i^{-1})\}^2$ and $E(R_i^{-3})/\{E(R_i^{-1})\}^3$. A straightforward approach is to consider moment estimators. We let $\hat{R}_i = \|X_i - \hat{\theta}_{n,p}\|$ and $\hat{R}_{i*} = \hat{R}_i + \hat{\theta}_{n,p}^\top \hat{U}_i - 2^{-1} \hat{R}_i^{-1} \|\hat{\theta}_{n,p}\|^2$. Then \hat{R}_{i*} can be viewed as a second-order approximation of R_i ; see Lemmas 1 and 2 in the Supplementary Material. Using \hat{R}_{i*} instead of \hat{R}_i would further reduce the bias in estimating $E(R_i^{-2})/E(R_i^{-1})^2$. Then the test statistic $(\tilde{Q} - p\hat{\delta}_{n,p})/\tilde{\sigma}_0$ converges to $N(0, 1)$ under H_0 as long as $np(\hat{\delta}_{n,p} - \delta_{n,p}) = o_p(1)$, where $\hat{\delta}_{n,p}$ is the estimator of $\delta_{n,p}$ upon replacing $E(R_i^{-k})/E(R_i^{-1})^k$ with $n^{k-1} \sum_{i=1}^n \hat{R}_{i*}^{-k} / (\sum_{i=1}^n \hat{R}_{i*}^{-1})^k$ in (5). The proposed test at significance level α rejects H_0 if $(\tilde{Q} - p\hat{\delta}_{n,p})/\tilde{\sigma}_0 > z_\alpha$, where z_α is the $1 - \alpha$ quantile of $N(0, 1)$.

Although Theorem 1 allows the dimensionality to increase at the rate of the square of the sample size, in practice how large p is allowed to be would depend mainly on the rate of ratio-consistency of $n \sum_{i=1}^n \hat{R}_{i*}^{-2} / (\sum_{i=1}^n \hat{R}_{i*}^{-1})^2$, to ensure that $np(\hat{\delta}_{n,p} - \delta_{n,p}) = o_p(1)$. Suppose that $n \sum_{i=1}^n \hat{R}_{i*}^{-2} / (\sum_{i=1}^n \hat{R}_{i*}^{-1})^2 = E(R_i^{-2})/E(R_i^{-1})^2 \{1 + O_p(a_n)\}$. It can be shown that under the null hypothesis and Assumption 1, $a_n = n^{-1/2}$. Also, $n^2 \sum_{i=1}^n \hat{R}_{i*}^{-3} / (\sum_{i=1}^n \hat{R}_{i*}^{-1})^3 \rightarrow E(R_i^{-3})/E(R_i^{-1})^3$ in probability. Thus, without imposing any other conditions, the bias-corrected method is valid when $p = o(n^{3/2})$. In

certain cases, a_n can be improved upon. For example, under the condition $\text{var}(R_i^{-k})/E(R_i^{-k})^2 = o(p^{-1/2})$ for $k = 1, 2$, we can get $a_n = o(n^{-1} + n^{-1/2}p^{-1/4})$. In such cases, $p = O(n^2)$ can be allowed. The multinormal distribution clearly satisfies this condition because we can show that $\text{var}(R_i^{-k})/E(R_i^{-k})^2 = O(p^{-1})$. The technical details can be found in the Supplementary Material.

Remark 1. When X_i comes from the normal distribution, $\lim_{p \rightarrow \infty} E(R_i^{-k})/E(R_i^{-1})^k = 1$ under the null hypothesis. In this case, $\delta_{n,p}$ can be simplified as $\delta_{n,p} \approx n^{-2} + 2n^{-3}$. We find that using this $\delta_{n,p}$ works almost as well as using $\hat{\delta}_{n,p}$ in all the cases considered; therefore it is recommended in practice when one wishes to reduce computational effort.

Next, we consider the asymptotic distribution of \tilde{Q} under the alternative hypothesis $H_1 : \Lambda_p = I_p + D_{n,p}$. Define

$$\tilde{\sigma}_1^2 = \tilde{\sigma}_0^2 + n^{-2}p^{-2} \{8p \text{tr}(D_{n,p}^2) + 4 \text{tr}^2(D_{n,p}^2)\} + 8n^{-1}p^{-2} \{\text{tr}(\Lambda_p^4) - p^{-1} \text{tr}^2(\Lambda_p^2)\}.$$

THEOREM 2. *Suppose that $n \text{tr}(D_{n,p}^2)/p = O(1)$ and $p = O(n^2)$. Under H_1 and Assumption 1, $\{\tilde{Q} - \text{tr}(D_{n,p}^2)/p - p\delta_{n,p}\}/\tilde{\sigma}_1 \rightarrow N(0, 1)$ in distribution as $p \rightarrow \infty$ and $n \rightarrow \infty$.*

A direct application of Theorem 2 is the consistency of the proposed test.

COROLLARY 1. *Suppose that the assumptions in Theorem 2 hold. If $n \text{tr}(D_{n,p}^2)/p \rightarrow \infty$, the test $\tilde{\sigma}_0^{-1}(\tilde{Q} - p\delta_{n,p}) > z_\alpha$ is consistent against H_1 as $p \rightarrow \infty$ and $n \rightarrow \infty$.*

Because $p^{-1} \text{tr}(D_{n,p}^2)$ measures the departure from the null hypothesis for the sphericity condition, this corollary ensures that as long as $p^{-1} \text{tr}(D_{n,p}^2)$ is not shrinking faster than n^{-1} , the tests are asymptotically optimal. The consistency rates implied in Corollary 1 by $n \text{tr}(D_{n,p}^2)/p \rightarrow \infty$ are, indeed, suboptimal in general. Neither John's test nor the sign-based test described here is asymptotically optimal, because their consistency rates are suboptimal. Under certain assumptions, such as multivariate normality, some rate-optimal tests can be constructed; see [Onatski et al. \(2013\)](#).

The following corollary provides the limiting efficiency comparison with the test of [Chen et al. \(2010\)](#) under multivariate normality. We consider the sequences of local alternatives $H_1 : \Lambda_p = I_p + D_{n,p}$ in which $C_1 \leq n \text{tr}(D_{n,p}^2)/p \leq C_2$ for two positive constants C_1 and C_2 .

COROLLARY 2. *Suppose that the assumptions in Theorem 2 hold. Under multinormal distributions, the sign-based test \tilde{Q} is asymptotically as efficient as the test of [Chen et al. \(2010\)](#).*

When the dimension p is fixed, it can be expected that the proposed test, which uses only the direction of an observation from the origin, will be outperformed by a test constructed with original observations, such as that of [Chen et al. \(2010\)](#). However, as $p \rightarrow \infty$ as $n \rightarrow \infty$, the disadvantage diminishes. It turns out that theoretically comparing the proposed test with the test of [Chen et al. \(2010\)](#) under general multivariate distributions is difficult. This is because the asymptotic validity of the test of [Chen et al. \(2010\)](#) relies on model (2), while an elliptical assumption is required in Theorems 1 and 2. Thus, in the next section, we compare these two methods using simulation.

3. NUMERICAL STUDIES

Three types of well-known multivariate elliptical distributions are considered: (I) the standard multivariate normal distribution; (II) the standard multivariate t distribution with four degrees of freedom, $t_{p,4}$; and (III) mixtures of two multivariate normal densities $\kappa f_p(\mu, I_p) + (1 - \kappa) f_p(\mu, 9I_p)$, where $f_p(\cdot, \cdot)$ is the p -variate multivariate normal density. The value κ is chosen to be 0.8 here. The model (2) is also included, to give us a broader picture of the robustness and efficiency of the proposed method. Following [Chen et al. \(2010\)](#), we chose $\Gamma = I_p$, and for each Z_i , p independent identically distributed random variables Z_{ij} were generated. Two distributions for the Z_{ij} are considered: (IV) the standardized gamma distribution $\text{Ga}(4, 0.5)$ considered by [Chen et al. \(2010\)](#); and (V) the standardized t distribution with four degrees of freedom, t_4 . The random vectors generated from scenarios IV and V are not elliptically distributed, while

Table 1. Empirical size and power (%) of four tests at the 5% significance level for multivariate normal random vectors, scenario I

	v	$p = 55$				$p = 181$				$p = 642$			
		T_{LW}	T_{CZZ}	T_{HP}	T_{BCS}	T_{LW}	T_{CZZ}	T_{HP}	T_{BCS}	T_{LW}	T_{CZZ}	T_{HP}	T_{BCS}
$n = 40$	0.000	4.8	5.6	17	4.9	4.8	5.8	74	4.9	5.8	6.2	100	5.1
	0.125	44	44	65	41	50	50	98	47	50	52	100	49
	0.250	68	69	84	64	72	72	99	68	74	74	100	72
$n = 80$	0.000	4.6	5.5	9.0	4.7	4.9	6.0	31	5.2	5.2	5.2	100	5.1
	0.125	87	87	90	84	93	92	99	91	95	95	100	94
	0.250	99	99	99	99	100	100	100	100	100	100	100	100

T_{LW} , the test of Ledoit & Wolf (2002); T_{CZZ} , the test of Chen et al. (2010); T_{HP} , the sign test (3) studied by Hallin & Paindaveine (2006); T_{BCS} , the proposed bias-corrected sign test.

Table 2. Empirical size and power (%) at the 5% significance level for multivariate t random vectors (scenario II) and mixtures of multivariate normal random vectors, scenario III

	v	Scenario II				Scenario III			
		$n = 40$		$n = 80$		$n = 40$		$n = 80$	
		T_{CZZ}	T_{BCS}	T_{CZZ}	T_{BCS}	T_{CZZ}	T_{BCS}	T_{CZZ}	T_{BCS}
$p = 55$	0.000	12	4.7	14	5.1	16	5.4	16	5.4
	0.125	34	39	61	84	37	40	57	82
	0.250	46	65	76	98	44	64	75	98
$p = 181$	0.000	12	5.9	14	4.4	15	5.0	16	4.5
	0.125	36	47	63	92	38	46	60	92
	0.250	47	69	79	99	44	69	79	100
$p = 642$	0.000	12	6.6	13	5.7	16	6.2	16	5.4
	0.125	35	50	61	94	35	50	62	94
	0.250	46	71	77	100	45	72	79	100

T_{CZZ} , the test of Chen et al. (2010); T_{BCS} , the proposed bias-corrected sign test.

neither scenario II nor scenario III corresponds to model (2); see the proof in the Supplementary Material. Only scenario I satisfies both the elliptical assumption and the form of model (2).

The combinations of p and n considered in Chen et al. (2009) are adopted here. The sample size n was taken to be 20, 40, 60 or 80, and dimensions $p = 38, 55, 89, 181, 331$ and 642 were considered for each sample size. Set $A = \text{diag}(2^{1/2}1_{[vp]}, 1_{p-[vp]})$, where $[x]$ denotes the integer truncation of x . To evaluate the size and power of the sphericity test, we generate multivariate random vectors, Y_i say, from scenarios I–V, and then obtain the observations $X_i = AY_i$. Three levels of v were considered: 0, 0.125 and 0.25. We compare the proposed test, hereafter called the bias-corrected sign test, with three methods for testing sphericity: the sign test (3) studied by Hallin & Paindaveine (2006), the test proposed by Ledoit & Wolf (2002), and the sphericity test proposed by Chen et al. (2010).

Tables 1 and 2 report the empirical size and power of the tests in scenarios I–III with $n = 40, 80$ and $p = 55, 181, 642$. The complete simulation results are given in the Supplementary Material. For each experiment we ran 2500 replications. The test of Ledoit & Wolf (2002) is not included in Table 2 because it is applicable only to normal distributions and encounters serious size distortion. Table 1 shows that the sign-based test without bias correction has Type I error rates much larger than the nominal levels, especially for large p and small n . This is consistent with the asymptotic analysis in § 2. For this reason, we do not report its results in Table 2.

Under scenario I, the empirical sizes of the tests converge to the nominal levels as p and n increase together. The test of Ledoit & Wolf (2002) performs best in terms of both size and power, as one would expect since it is based on normality. The bias-corrected sign test has similar empirical sizes. The power of the proposed test is largely dependent on the sample size and levels of v , as these determine $\text{tr}(D_{n,p}^2)$.

The test of [Chen et al. \(2010\)](#) outperforms the bias-corrected sign test in terms of power in most cases, but as n and p increase the advantage tends to diminish. When $n = 80$ and $p \geq 181$, the two tests are largely comparable and the difference is at least partly due to the unequal sizes of the tests. This can be understood from Corollary 2.

Table 2 presents values obtained from simulations with the other two elliptical distributions. The proposed test can achieve the nominal size, whereas the test of [Chen et al. \(2010\)](#) has considerable bias in size; even worse, the empirical size of the latter test hardly improves as n and/or p increase. In both scenarios the bias-corrected sign test is more efficient under H_1 in the sense that even when the observed size is much smaller than that of [Chen et al. \(2010\)](#), the empirical power increases much faster as v increases. When $n = 80$, the proposed test performs uniformly much better than the test of [Chen et al. \(2010\)](#), and the difference is quite remarkable. Certainly, this is not surprising as neither $t_{p,4}$ nor a mixture of multivariate normal distributions belongs to model (2), on which the validity of the test of [Chen et al. \(2010\)](#) depends so much.

The empirical size and power of the tests in scenarios IV and V are reported in the Supplementary Material. Although our test is not asymptotically justified under model (2), it is quite robust with respect to the two distributions which belong to that model. Its sizes are close to nominal and even closer than those of the test of [Chen et al. \(2010\)](#). With regard to performance under alternatives, our test has quite good power and generally performs similarly to the test of [Chen et al. \(2010\)](#), although its sizes are usually smaller. These results suggest that the proposed test is quite robust and efficient in testing sphericity, especially for heavy-tailed or skewed distributions.

4. CONCLUDING REMARKS

The bias-correction procedure takes advantage of the relatively simple form of multivariate sign-based tests for sphericity. We believe that this procedure can be extended to more general elliptical distributions with $\Sigma_p = \text{diag}\{\sigma_{11}, \dots, \sigma_{pp}\}$ where the σ_{ii} are unknown. Moreover, Theorem 1 is established under $p = O(n^2)$. The issue preventing p from growing faster than n^2 is that a higher-order expansion is required for bias correction.

[Hallin & Paindaveine \(2006\)](#) proposed a family of signed-rank test statistics based on the sign vectors U_i and the ranks of the moduli R_i . Their tests appear to be asymptotically optimal at given target densities. Deriving similar bias-corrected procedures for those tests is difficult because of their complicated construction and merits further research. The tests based on symmetrized spatial-signs and spatial-ranks derived in [Sirkiä et al. \(2009\)](#) also warrant future study in a high-dimensional setting.

SUPPLEMENTARY MATERIAL

Supplementary material available at *Biometrika* online includes the proofs of Theorems 1 and 2, the calculation of $E(\Delta_i)$, additional simulation results and some other technical details.

ACKNOWLEDGEMENT

The authors would like to thank the editor, associate editor and two anonymous referees for their many helpful comments that have improved the article significantly. This research was supported by the National Natural Science Foundation and Research Fund for the Doctoral Program of Higher Education of China, the Foundation for the Authors of National Excellent Doctoral Dissertations, and the Program for New Century Excellent Talents in University.

REFERENCES

- BARINGHAUS, L. (1991). Testing for spherical symmetry of a multivariate distribution. *Ann. Statist.* **19**, 899–917.
 CHEN, S. X., PENG, L. & QIN, Y. L. (2009). Effects of data dimension on empirical likelihood. *Biometrika* **96**, 1–12.

- CHEN, S. X. & QIN, Y. L. (2010). A two-sample test for high-dimensional data with applications to gene-set testing. *Ann. Statist.* **38**, 808–35.
- CHEN, S. X., ZHANG, L. X. & ZHONG, P. S. (2010). Tests for high-dimensional covariance matrices. *J. Am. Statist. Assoc.* **105**, 801–15.
- GHOSH, S. K. & SENGUPTA, D. (2001). Testing for proportionality of multivariate dispersion structures using interdirections. *J. Nonparam. Statist.* **13**, 331–49.
- HALLIN, M. & PAINDAVEINE, D. (2006). Semiparametrically efficient rank-based inference for shape. I. optimal rank-based tests for sphericity. *Ann. Statist.* **34**, 2707–56.
- HJORT, H. L., MCKEAGUE, I. W. & VAN KEILEGOM, I. (2009). Extending the scope of empirical likelihood. *Ann. Statist.* **37**, 1079–115.
- JOHN, S. (1971). Some optimal multivariate tests. *Biometrika* **59**, 123–7.
- JOHN, S. (1972). The distribution of a statistic used for testing sphericity of normal distributions. *Biometrika* **59**, 169–73.
- LEDOIT, O. & WOLF, M. (2002). Some hypothesis tests for the covariance matrix when the dimension is large compared to the sample size. *Ann. Statist.* **30**, 1081–102.
- MARDEN, J. & GAO, Y. (2002). Rank-based procedures for structural hypotheses on covariance matrices. *Sankhyā Ser. A* **64**, 653–77.
- MAUCHLY, J. W. (1940). Significance test for sphericity of a normal n -variate distribution. *Ann. Math. Statist.* **11**, 204–9.
- MÖTTÖNEN J. & OJA, H. (1995). Multivariate spatial sign and rank methods. *J. Nonparam. Statist.* **5**, 201–13.
- MUIRHEAD, R. J. & WATERNAUX, C. M. (1980). Asymptotic distributions in canonical correlation analysis and other multivariate procedures for nonnormal populations. *Biometrika* **67**, 31–43.
- OJA, H. (2010). *Multivariate Nonparametric Methods with R*. New York: Springer.
- ONATSKI, A., MOREIRA, M. J. & HALLIN, M. (2013). Asymptotic power of sphericity tests for high-dimensional data. *Ann. Statist.* **41**, 1204–31.
- SCHOTT, J. R. (2005). Testing for complete independence in high dimensions. *Biometrika* **92**, 951–6.
- SIRKIÄ, S., TASKINEN, S., OJA, H. & TYLER, D. E. (2009). Tests and estimates of shape based on spatial signs and ranks. *J. Nonparam. Statist.* **21**, 155–76.
- TANG, C. Y. & LENG, C. (2010). Penalized high-dimensional empirical likelihood. *Biometrika* **97**, 905–20.
- TYLER, D. E. (1987). Statistical analysis for the angular central Gaussian distribution on the sphere. *Biometrika* **74**, 579–89.

[Received March 2012. Revised August 2013]